arXiv:math/0107189v2 [math.AG] 3 Jul 2003

LOCAL ZETA FUNCTION FOR CURVES, NON DEGENERACY CONDITIONS AND NEWTON POLYGONS

M.J. SAIA AND W. A. ZUNIGA-GALINDO

Abstract. This paper is dedicated to the description of the poles of the Igusa local zeta function $Z(s,f,v)$ when $f(x,y)$ satisfies a new non degeneracy condition called *arithmetic non degeneracy.* More precisely, we attach to each polynomial $f(x,y)$ a collection of convex sets $\Gamma^{A}(f)=\{\Gamma_{f,1},\cdots,\Gamma_{f,l_0}\}$ called the *arithmetic Newton polygon* of $f(x,y)$, and introduce the notion of *arithmetic non degeneracy with respect to* $\Gamma^{A}(f)$. If L_v is a $p-$adic field, and $f(x,y)\in L_v\left[x,y\right]$ is arithmetically non degenerate, then the poles of $Z(s,f,v)$ can be described explicitly in terms of the equations of the straight segments that conform the boundaries of the convex sets $\Gamma_{f,1},\cdots,\ \Gamma_{f,l_0}$. Moreover, the proof of main result gives an effective procedure for computing $Z(s,f,v)$.

1. Introduction

Let L_v be a $p-$adic field, and v the corresponding a non-archimedean valuation. Let O_v the ring of integers of L_v, P_v the maximal ideal of O_v, and $\overline{L_v}=O_v/\ P_v$ the residue field of L_v. The cardinality of $\overline{L_v}$ is denoted by q, thus $\overline{L_v}=\mathbb{F}_q$. We fix a local parameter π for O_v. For $z\in L_v$, $\mid z\mid_v:=q^{-v(z)}$ denotes its normalized absolute value. Given a non-constant polynomial $g(x,y)\in L_v[x,y]$, the Igusa local zeta function $Z(s,g,v)$ attached to $g(x,y)$ and L_v is defined as

$$Z(s,g,v):=\int\limits_{O_v^2}\mid g(x,y)\mid_v^s\mid dxdy\mid,$$

where $s\in\mathbb{C}$ satisfying $Re(s)>0$ and $\mid dxdy\mid$ denotes a Haar measure of L_v^2 normalized so that the measure of O_v^2 is one.

Igusa showed that the local zeta function $Z(s,g,v)$ is a rational function of q^{-s}, for general polynomials with coefficients in a $p-$adic

---

1991 *Mathematics Subject Classification.* Primary 11D79, 14G20, 14M25.

Key words and phrases. Igusa local zeta functions, Newton polygons, degenerate curves, non degeneracy conditions, polynomial congruences.

The first named author was partially supported by CNPq-Grant 300556/92-6.

The second named author was supported by COLCIENCIAS-Grant # 089-2000. The second named author also thanks the partial support given by FAPESP for visiting the Instituto de Matemática e Computaçao, Universidade de São Paulo, Campus São Carlos, in January 2000.

field [5]. Igusa's local zeta functions are related to the number of solutions of congruences modulo p^m and to exponential sums modulo p^m. There are several conjectures and intriguing connections between Igusa's local zeta functions with topology and singularity theory [2].

This paper is dedicated to the description of the poles of the Igusa local zeta functions $Z(s, f, v)$ when $f(x, y)$ satisfies a new non degeneracy condition called *arithmetic non degeneracy*. More precisely, we attach to each polynomial $f(x, y)$ a collection of convex sets $\Gamma^A(f) = \{\Gamma_{f,1}, .., \Gamma_{f,l_0}\}$ called the *arithmetic Newton polygon* of $f(x, y)$, and introduce the notion of *arithmetic non degeneracy with respect to* $\Gamma^A(f)$ (see section 4).

The set of degenerate polynomials, in the usual sense, is a proper subset of the set of arithmetically non degenerate polynomials. In addition, for a fixed geometric Newton polygon $\Gamma^{geom} \subset \mathbb{R}^2$, the set of degenerate complex polynomials in two variables with respect Γ^{geom} contains an open set, for the Zariski topology, consisting of arithmetically non degenerate polynomials.

Our main result gives an explicit list for the possible poles of $Z(s, f, v)$, when $f(x, y) \in L_v[x, y]$ is an arithmetically non degenerate polynomial, in terms of the equations of the straight segments that conform the boundaries of the convex sets $\Gamma_{f,1}$,.., Γ_{f,l_0} (see theorems 5.1, 6.1). Moreover, our proof gives an effective procedure for computing $Z(s, f, v)$.

The notion of non degeneracy due Kouchnirenko [9] is useful in the computation of Milnor numbers, number of isolated solutions of polynomial equations, etc. The authors hope that the notion of arithmetic non degeneracy may be useful for other purposes, like those above mentioned, different from the description of the poles of local zeta functions.

The description of the poles of the local zeta functions in terms of Newton polygons is a well known fact for non degenerate polynomials in the sense of Kouchnirenko [13], [10], [3], [4], [16], [17]. In the degenerate case there are no previous results showing a relation between the poles of local zeta functions and Newton polygons.

The poles of $Z(s, f, v)$, for a general polynomial f in two variables, can be described explicitly in terms of a resolution of singularities for f (see [14], [15], [11], [7], [12]). The main result of this paper is complementary to the mentioned results. Indeed, our main result produces an explicit list of candidates for the poles of $Z(s, f, v)$ without using resolutions of singularities, but this result is only valid for a large class of polynomials.

In this paper, we work with $p-$adic fields of characteristic zero, however all results are valid in positive characteristic.

This paper is organized as follows. In the section 2, we review some well-known results about geometric Newton polygons, non degeneracy conditions in the sense of Kouchnirenko, and certain $p-$adic integrals attached to non degenerate polynomials. In section 3, we compute explicitly the local zeta function of some degenerate polynomials, in the sense of Kouchnirenko. These examples constitute the basic models for the effective computation of the local zeta functions of arithmetically non degenerate polynomials. In section 4, we introduce the notion of arithmetic Newton polygon and arithmetic non degeneracy condition. In section 5, we compute the several $p-$adic integrals that appear in the effective computation of the local zeta function of an arithmetically non degenerate polynomial. In section 6, we prove the main result.

2. Preliminaries

2.1. **Geometric Newton polygons.**

We set $\mathbb{R}_+ = \{x \in \mathbb{R} \mid x \geqq 0\}$. Let $f(x,y) = \sum_{(i,j)} a_{i,j}x^i y^j$ be a non-constant polynomial in two variables with coefficients in a field K satisfying $f(0,0) = 0$. The set *supp*$(f) = \{(i,j) \in \mathbb{N}^2 \mid a_{i,j} \neq 0\}$ is called the *support* of f. *The geometric Newton polygon* $\Gamma^{geom}(f)$ of f is defined as the convex hull in $\mathbb{R}^2_+$ of the set $\cup_{(i,j)\in supp(f)} \left((i,j) + \mathbb{R}^2_+\right)$.

By a *proper face* γ of $\Gamma^{geom}(f)$, we mean a non-empty convex set γ obtained by intersecting $\Gamma^{geom}(f)$ with an straight line H, such that $\Gamma^{geom}(f)$ is contained in one of two half-spaces determined by H. The straight line H is called *the supporting line* of γ. Thus the faces can be points (zero-dimensional faces) or straight segments (one-dimensional faces). The last ones are also called *facets*.

If γ is a face of $\Gamma^{geom}(f)$ its *face function* is defined to be the polynomial

$$f_\gamma(x,y) := \sum_{(i,j)\in\gamma} a_{i,j}x^i y^j.$$

Definition 2.1. *A non-constant polynomial $f(x,y)$ satisfying $f(0,0) = 0$ is called non degenerate with respect to $\Gamma^{geom}(f)$, in the sense Kouchnirenko, if it satisfies:*

1. *the origin of K^2 is a singular point of $f(x,y)$;*
2. *for each face $\gamma \subseteq \Gamma^{geom}(f)$, including $\Gamma^{geom}(f)$ itself, the system of equations*
$$f_\gamma(x,y) = \frac{\partial f_\gamma}{\partial x}(x,y) = \frac{\partial f_\gamma}{\partial y}(x,y) = 0,$$
does not have solutions on $K^{\times 2}$.

We set $\langle .,. \rangle$ for the usual inner product in $\mathbb{R}^2$, and identify the dual vector space with $\mathbb{R}^2$. For $a \in \mathbb{R}^2_+$, we define

$$m(a) := \inf_{x\in\Gamma(f)} \{\langle a, x\rangle\}.$$

Given any $a \in \mathbb{R}^2_+ \smallsetminus \{0\}$ *the first meet locus* $F(a)$ of a is defined as

$$F(a) := \{x \in \Gamma^{geom}(f) \mid \langle a, x\rangle = m(a)\}.$$

The first meet locus $F(a)$ of $a \in \mathbb{R}_+^2 \smallsetminus \{0\}$ is a proper face of $\Gamma^{geom}(f)$.

Given a proper face γ of $\Gamma^{geom}(f)$, the *cone associated* to γ is defined to be the set

$$\Delta_\gamma := \left\{ a \in \mathbb{R}_+^2 \smallsetminus \{0\} \mid F(a) = \gamma \right\}.$$

We define an equivalence relation on $\mathbb{R}_+^2 \smallsetminus \{0\}$ by

$$a \backsimeq a' \text{ if and only if } F(a) = F(a').$$

The equivalence classes of $\backsimeq$ are the cones Δ_γ. The following two propositions describe the geometry of the equivalences classes of $\backsimeq$ (see e.g. [13], [3]).

Proposition 2.1. *Let γ be a proper face of $\Gamma^{geom}(f)$. If γ is a one-dimensional face and a is perpendicular vector on γ, then*

$$\Delta_\gamma = \{\alpha a \mid \alpha \in \mathbb{R}, \ \alpha > 0\}. \tag{2.1}$$

If γ is a zero-dimensional face and w_1, w_2 are the facets of $\Gamma^{geom}(f)$ which contain γ, and a_1, a_2 are vectors which are perpendicular on respectively w_1, w_2, then

$$\Delta_\gamma = \{\alpha_1 a_1 + \alpha_2 a_2 \mid \alpha_i \in \mathbb{R}, \ \alpha_i > 0\}. \tag{2.2}$$

A set of the form (2.2) (respectively (2.1)) is called a *cone strictly positive spanned* by the vectors $\{a_1, a_2\}$ (respectively by $\{a\}$). Let Δ be a strictly positive spanned by a vector set A, with $A = \{a_1, a_2\}$, or $A = \{a\}$. If A is linearly independent over $\mathbb{R}$, the cone Δ is called *a simplicial cone.* In this last case, if the elements of A are in $\mathbb{Z}^2$, the cone Δ is called a *rational simplicial* cone. If A can be completed to be a basis of $\mathbb{Z}-$module $\mathbb{Z}^2$, the cone Δ is called *a simple cone.*

A vector $a \in \mathbb{R}^2$ is called *primitive* if the components of a are positive integers whose greatest common divisor is one.

For every facet of $\Gamma^{geom}(f)$ there is a unique primitive vector in $\mathbb{R}^2$ which is perpendicular to this facet. We denote by D the set of such vectors. Thus each equivalence class under $\backsimeq$ is a rational simplicial cone spanned by elements of D.

From above considerations follow that there exists a partition of $\mathbb{R}_+^2$ of the form:

$$\mathbb{R}_+^2 = \{(0,0)\} \bigcup \bigcup_{\gamma \subset \Gamma^{geom}(f)} \Delta_\gamma, \tag{2.3}$$

where γ runs through all proper faces of $\Gamma^{geom}(f)$. We shall say that $\{\Delta_\gamma\}_{\gamma \subset \Gamma^{geom}(f)}$ is a *simplicial conical subdivision* of $\mathbb{R}_+^2$ subordinated to $\Gamma^{geom}(f)$.

2.2. **Local zeta functions and conical subdivisions.** Let $f(x,y) \in L_v[x,y]$ be a non-constant polynomial satisfying $f(0,0)=0$, and $\Gamma^{geom}(f)$ its geometric Newton polygon. We fix a simplicial conical subdivision $\{\Delta_\gamma\}_{\gamma \subset \Gamma^{geom}(f)}$ of $\mathbb{R}_+^2$ subordinated to $\Gamma^{geom}(f)$. We define

$$E_{\Delta_\gamma} := \{(x,y) \in O_v^2 \mid (v(x), v(y)) \in \Delta_\gamma\},$$

$$Z(s,f,v,\Delta_\gamma) := \int\limits_{E_{\Delta_\gamma}} \mid f(x,y) \mid_v^s \mid dxdy \mid,$$

and

$$Z(s,f,v,O_v^{\times 2}) := \int\limits_{O_v^{\times 2}} \mid f(x,y) \mid_v^s \mid dxdy \mid .$$

With the above notation, it follows from partition (2.3) that

$$Z(s,f,v) = Z(s,f,v,O_v^{\times 2}) + \sum_{\gamma \subset \Gamma^{geom}(f)} Z(s,f,v,\Delta_\gamma). \tag{2.4}$$

Therefore the computation of the integral $Z(s,f,v)$ is reduced to the computation of integrals of type $Z(s,f,v,O_v^{\times 2})$ and $Z(s,f,v,\Delta_\gamma)$. In the case in which f is non degenerate with respect $\Gamma^{geom}(f)$, the above mentioned integrals can be computed explicitly using techniques of toroidal geometry, and the $p-$adic stationary phase formula [5, theorem 10.2.1], [4], [16], [17], [18].

For a general f formula (2.4) is still valuable, if f does not have singularities on $(L_v^\times)^2$. By applying the stationary phase formula (abbreviated SPF) repeatedly, it is possible to compute effectively $Z(s,f,v,O_v^{\times 2})$ (see [16, lemma 3.1], or [17, lemma 2.4]). For a future reference we restate this result here.

Lemma 2.1. *Let $g_m(x,y) = g_0(x,y) + \pi^m g_1(x,y) \in L_v[x,y]$ be a non-constant polynomial, $m \geqslant 1$, such that $g_m(x,y)$,and $g_0(x,y)$ do not have singular points on $(L_v^\times)^2$. There exists a constant $c(g_0)$ such that if $m \geqq c(g_0)$, then*

$$Z(s,g_m,v,O_v^{\times 2}) = \int\limits_{O_v^{\times 2}} \mid g_0(x,y) \mid_v^s \mid dxdy \mid = \frac{U(q^{-s})}{1-q^{-1-s}},$$

with $U(q^{-s}) \in \mathbb{Q}[q^{-s}]$.

The computation of the integrals $Z(s,f,v,\Delta_\gamma)$ presents two cases. If f_γ does not have singularities on $(L_v^\times)^2$, we shall say that f *is non degenerate on* γ. In this case the integrals $Z(s,f,v,\Delta_\gamma)$ can be computed by using known techniques [10], [4], [16], [17]. For a future reference we restate this result here.

Lemma 2.2. *Let $f(x,y) \in L_v[x,y]$ be a non-constant polynomial such that $f(0,0)=0$, and f does not have singular points on $(L_v^\times)^2$.*

(1) *If* Δ_γ *is a two-dimensional cone, and* $f_\gamma(x,y)$ *does not have singularities on the torus* $(L_v^\times)^2$*, then*

$$Z(s,f,v,\Delta_\gamma) = U_1(q^{-s}) \in \mathbb{Q}[q^{-s}].$$

(2) *If* Δ_γ *is a one-dimensional cone, and* $f_\gamma(x,y)$ *does not have singularities on* $(L_v^\times)^2$*, then*

$$Z(s,f,v,\Delta_\gamma) = \frac{U_2(q^{-s})}{(1-q^{-1-s})(1-q^{-(a+b)-d_\gamma s})},\ U_2(q^{-s}) \in \mathbb{Q}[q^{-s}],$$

where $ax+by=d_\gamma$ *is the equation of the supporting line of the facet* γ*.*

The core of this paper is the explicit computation of integrals of type

$$Z(s,f,v,\Delta_\gamma),$$

when f is *degenerate on* Δ_γ, but satisfies a new non degeneracy condition. This will be done in sections 4, 5.

Remark 2.1. *Given a* Δ *be a rational simplicial cone, there exists a partition of* Δ *into simple cones* [8, page 32-33]*. This fact can be used to obtain a partition of* $\mathbb{R}_+^2$ *in simple cones. In this case, it is possible to compute the integral* $Z(s,f,v,\Delta)$ *in a simple form, when* f *is non degenerate, in the sense of Kouchnirenko. However this procedure produces a larger list of candidates to poles for* $Z(s,f,v,\Delta)$*.*

3. Model cases

In this section we shall compute explicitly the local zeta functions for some degenerate polynomials, in the sense of Kouchnirenko. These examples constitute the basic models for the effective computation of the local zeta functions of arithmetically non degenerate polynomials.

3.1. **The local zeta function of** $(y^3-x^2)^2+x^4y^4$**.**

We assume that the characteristic of the residue field of L_v is different from 2. The origin of L_v^2 is the only singular point of $f(x,y)=(y^3-x^2)^2+x^4y^4$.This polynomial is degenerate with respect to its geometric Newton polygon.

The conical subdivision of $\mathbb{R}_+^2$ subordinated to the geometric Newton polygon of $f(x,y)$ is

$$\mathbb{R}_+^2 = \{(0,0)\} \bigcup \bigcup_{j=1}^{9} \Delta_j, \tag{3.1}$$

with

(3.2)

$\Delta_1 := (0,1)\mathbb{R}_+$
$\Delta_2 := (0,1)\mathbb{R}_+ + (1,1)\mathbb{R}_+$
$\Delta_3 := (1,1)\mathbb{R}_+$
$\Delta_4 := (1,1)\mathbb{R}_+ + (3,2)\mathbb{R}_+$
$\Delta_5 =: (3,2)\mathbb{R}_+$
$\Delta_6 := (3,2)\mathbb{R}_+ + (2,1)\mathbb{R}_+$
$\Delta_7 := (2,1)\mathbb{R}_+$
$\Delta_8 := (2,1)\mathbb{R}_+ + (1,0)\mathbb{R}_+$
$\Delta_9 := (1,0)\mathbb{R}_+$

In addition, all the Δ_i are rational simple cones. By using the above subdivision is possible to reduce the computation of $Z(s,f,v)$ to the computation of $Z(s,f,v,O_v^{\times 2})$, and $Z(s,f,v,\Delta_i)$, $i = 1,2,3,4,6,7,8,9$ (see2.2).

3.2. **Computation of** $Z(s,f,v,O_v^{\times 2})$**.** Since $f(x,y)$ does not have singularities on $(L_v^{\times})^2$ the integral can be computed by using SPF [5, theorem 10.2.1]:

$$Z(s,f,v,O_v^{\times 2}) = \left[\left(1-q^{-1}\right)^2 - q^{-2}N(f)\right] + \frac{q^{-2}(N(f)-1)\,(1-q^{-1})\,q^{-s}}{1-q^{-1-s}},$$

with $N(f) = Card\left(\left\{(u,v)\in\left(\mathbb{F}_q^{\times}\right)^2 \mid f(u,v)=0\right\}\right)$.

3.3. **Computation of** $Z(s,f,v,\Delta_i)$, $i = 1,2,3,4,6,7,8,9$**.** These integrals correspond to the case in which f is non degenerate on Δ_i, $i =$ 1, 2, 3, 4, 6, 7, 8, 9. The integrals corresponding to Δ_i, $i = 1,2$, can be calculated as follows.

3.3.1. *Case* $Z(s,f,v,\Delta_1)$.

$$\begin{aligned}
Z(s,f,v,\Delta_1) &= \sum_{n=1}^{\infty}\int\limits_{O_v^{\times}\times\pi^n O_v^{\times}} \mid f(x,y)\mid_v^s \mid dxdy \mid \\
&= \sum_{n=1}^{\infty} q^{-n}\int\limits_{O_v^{\times 2}} \mid (\pi^{3n}y^3 - x^2)^2 + \pi^{4n}x^4y^4 \mid_v^s \mid dxdy \mid \\
&= (1-q^{-1})^2\sum_{n=1}^{\infty} q^{-n} = q^{-1}(1-q^{-1}).
\end{aligned}$$

3.3.2. *Case* $Z(s,f,v,\Delta_2)$.

$$Z(s,f,v,\Delta_2) = \sum_{m=1}^{\infty}\sum_{n=1}^{\infty}\int\limits_{\pi^m O_v^{\times}\times\pi^{n+m}O_v^{\times}} \mid f(x,y)\mid_v^s \mid dxdy \mid =$$

$$\sum_{m=1}^{\infty}\sum_{n=1}^{\infty} q^{-2m-n-4ms}\int\limits_{O_v^{\times 2}} \mid (\pi^{3n+m}y^3-x^2)^2+\pi^{4m+4n}x^4y^4\mid_v^s \mid dxdy \mid$$

$$= (1-q^{-1})q^{-1}\frac{q^{-2-4s}}{(1-q^{-2-4s})}.$$

The other integrals can be computed in the same form. The following table summarizes these computations:

(3.3)

Cone	$Z(s,f,v,\Delta)$
Δ_1	$q^{-1}(1-q^{-1})$
Δ_2	$(1-q^{-1})q^{-1}\frac{q^{-2-4s}}{(1-q^{-2-4s})}$
Δ_3	$(1-q^{-1})^2\frac{q^{-2-4s}}{(1-q^{-2-4s})}$
Δ_4	$(1-q^{-1})^2\frac{q^{-7-16s}}{(1-q^{-2-4s})(1-q^{-5-12s})}$
Δ_6	$\frac{(1-q^{-1})^2q^{-8-18s}}{(1-q^{-5-12s})(1-q^{-3-6s})}$
Δ_7	$\frac{(1-q^{-1})^2q^{-3-6s}}{(1-q^{-3-6s})}$
Δ_8	$\frac{(1-q^{-1})q^{-4-6s}}{(1-q^{-3-6s})}$
Δ_9	$q^{-1}(1-q^{-1})$

3.4. **Computation of $Z(s,f,v,\Delta_5)$ (an integral on a degenerate face in the sense of Kouchnirenko).**

$$Z(s,f,v,\Delta_5) = \sum_{n=1}^{\infty}\int\limits_{\pi^{3n}O_v^{\times}\times\pi^{2n}O_v^{\times}} \mid f(x,y)\mid_v^s \mid dxdy \mid$$

$$= \sum_{n=1}^{\infty} q^{-5n-12ns}\int\limits_{O_v^{\times 2}} \mid (y^3-x^2)^2+\pi^{8n}x^4y^4\mid_v^s \mid dxdy \mid . \tag{3.4}$$

We set $f^{(n)}(x,y) = (y^3-x^2)^2+\pi^{8n}x^4y^4$, for $n \geqslant 1$. In order to compute (3.4), we first compute the integral

$$I(s,f^{(n)},v) := \int\limits_{O_v^{\times 2}} \mid (y^3-x^2)^2+\pi^{8n}x^4y^4\mid_v^s \mid dxdy \mid ,\ n \geqslant 1. \tag{3.5}$$

For that, we use the following change of variables

$$\begin{array}{rccc} \Phi : & O_v^{\times 2} & \longrightarrow & O_v^{\times 2} \\ & (x,y) & \longrightarrow & (x^3y, x^2y). \end{array} \tag{3.6}$$

The map Φ gives an analytic bijection of $O_v^{\times 2}$ onto itself, and preserves the Haar measure because its Jacobian $J_\Phi(x,y) = x^4 y$ satisfies $\mid J_\Phi(x,y) \mid_v = 1$, for every $x, y \in O_v^\times$. Thus

$$f^{(n)} \circ \Phi(x,y) = x^{12} y^4 \widetilde{f^{(n)}}(x,y), \tag{3.7}$$

with

$$\widetilde{f^{(n)}}(x,y) := (y-1)^2 + \pi^{8n} x^8 y^4, \tag{3.8}$$

and then from (3.6), and (3.7), it follows that

$$I(s, f^{(n)}, v) = \int\limits_{O_v^{\times 2}} \mid \widetilde{f^{(n)}}(x,y) \mid_v^s \mid dxdy \mid . \tag{3.9}$$

In order to compute the integral $I(s, f^{(n)}, v)$, $n \geqslant 1$, we decompose $O_v^{\times 2}$ as follows

$$O_v^{\times 2} = \bigcup_{y_0 \not\equiv 1 \bmod \pi} O_v^\times \times \{y_0 + \pi O_v\} \bigcup O_v^\times \times \{1 + \pi O_v\}, \tag{3.10}$$

where y_0 runs through a set of representatives of $\mathbb{F}_q^\times$ in O_v. From partition (3.10), and (3.8), it follows that

(3.11)

$$I(s, f^{(n)}, v) = (q-2) q^{-1} (1 - q^{-1}) + \int\limits_{O_v^\times \times \{1 + \pi O_v\}} \mid \widetilde{f^{(n)}}(x,y) \mid_v^s \mid dxdy \mid .$$

The integral in the right side of (3.11), admits the following expansion:

(3.12)

$$I(s, f^{(n)}, v) = (q-2) q^{-1} (1 - q^{-1}) + \sum_{j=1}^{\infty} q^{-j} \int\limits_{O_v^{\times 2}} \mid \widetilde{f^{(n)}}(x, 1 + \pi^j z) \mid_v^s \mid dxdz \mid,$$

where $\widetilde{f^{(n)}}(x, 1 + \pi^j z) = \pi^{2j} z^2 + \pi^{8n} x^8 (1 + \pi^j z)^4$. In the next step, we compute explicitly $\mid \widetilde{f^{(n)}}(x, 1 + \pi^j z) \mid_v^s$, for a fixed $n \geqq 1$. In this case, this can be done by a simple calculation:

(3.13)

$$\mid \widetilde{f^{(n)}}(x, 1 + \pi^j z) \mid_v^s = \begin{cases} q^{-2js} & \text{if} \quad 1 \leqq j \leqq 4n - 1, \\ q^{-8ns} \mid z^2 + x^8 (1 + \pi^j z)^4 \mid_v^s & \text{if} \quad j = 4n, \\ q^{-8ns} & \text{if} \quad j \geqq 4n + 1, \end{cases}$$

for any $(x, z) \in O_v^{\times 2}$.

We note that the explicit computation of the absolute value of $\mid \widetilde{f^{(n)}}(x, 1 + \pi^j z) \mid_v$ depends on an explicit description of the set $\{(w, z) \in \mathbb{R}^2 \mid w \leqq \min\{2z, 8n\}\}$, for a fixed n.

Now from (3.12), (3.13) , and the identity $\sum_{k=A}^{B} z^k = \frac{z^A - z^{B+1}}{1-z}$, we obtain an explicit expression for $I(s, f_n, v)$:

$$\begin{aligned} I(s, f^{(n)}, v) &= (q-2)q^{-1}(1-q^{-1}) + (1-q^{-1})^2 \frac{q^{-1-2s}}{(1-q^{-1-2s})} \\ &\quad -(1-q^{-1})^2 \frac{q^{-4n-8ns}}{(1-q^{-1-2s})} + q^{-4n-8ns} I_0(s) \\ &\quad +q^{-1}(1-q^{-1})q^{-4n-8ns}, \end{aligned} \tag{3.14}$$

where

$$I_0(s) := \int\limits_{O_v^{\times 2}} \mid z^2 + x^8(1+\pi^{4n}z)^4 \mid_v^s \mid dxdz \mid . \tag{3.15}$$

Since the reduction modulo π of the polynomial $z^2 + x^8(1+\pi^{4n}z)^4$ does not have singular points on $\mathbb{F}_q^{\times 2}$, because the characteristic of the residue field is different of 2, SPF implies that $I_0(s) = \frac{U_1(q^{-s})}{(1-q^{-1-s})}$, where $U_1(q^{-s})$ is a polynomial in q^{-s} independent of j and n.

Finally, from (3.4), (3.9) and (3.14), we obtain the following explicit expression for $Z(s, f, v, \Delta_5)$:

$$\begin{aligned} Z(s, f, v, \Delta_5) &= \frac{(q-2)(1-q^{-1})q^{-6-12s}}{(1-q^{-5-12s})} + \frac{(1-q^{-1})^2 q^{-6-14s}}{(1-q^{-1-2s})(1-q^{-5-12s})} \\ &\quad -\frac{(1-q^{-1})^2 q^{-9-20s}}{(1-q^{-1-2s})(1-q^{-9-20s})} + \frac{q^{-9-20s}U_1(q^{-s})}{(1-q^{-1-s})(1-q^{-9-20s})} \\ &\quad +\frac{(1-q^{-1})q^{-10-20s}}{(1-q^{-9-20s})}. \end{aligned} \tag{3.16}$$

From the explicit formulas for $Z(s, f, v, \Delta_i)$, $i = 1, 2, \cdots, 8$, and $Z(s, f, v, O_v^{\times 2})$ (see (3.3), (3.16)), it follows that the real parts of the poles of $Z(s, f, v)$ belong to the set

$$\{-1\} \bigcup \{-\frac{5}{12}\} \bigcup \{-\frac{1}{2}, -\frac{9}{20}\}. \tag{3.17}$$

3.5. **The local zeta function of** $(y^3 - x^2)^2(y^3 - ax^2) + x^4y^4$**.** We set $g(x, y) = (y^3 - x^2)^2(y^3 - ax^2) + x^4y^4 \in L_v[x, y]$, with $a \in O_v^{\times}$, and $a \not\equiv 1 \bmod \pi$. We assume that the characteristic of the residue field of L_v is different from 2. The polynomial g is degenerate with respect to its geometric Newton polygon. In addition, the origin of L_v^2 is the only singular point of g.

By using the same decomposition of $\mathbb{R}_+^2$ into rational simple cones as in the above example, it is possible to reduce the computation of $Z(s, g, v)$ to the computation of the $p-$adic integrals $Z(s, g, v, O_v^{\times 2})$, $Z(s, g, v, \Delta_i), i = 1, \cdots, 8, 9$.

3.6. **Computation of $Z(s,g,v,O_v^{\times 2})$.** Since $g(x,y)$ does not have singularities on $(L_v^{\times})^2$ the integral can be computed by using SPF [5, theorem 10.2.1]:

$$Z(s,g,v,O_v^{\times 2}) = \left[\left(1-q^{-1}\right)^2 - q^{-2}N(g)\right] + \frac{q^{-2}(N(g)-1)\left(1-q^{-1}\right)q^{-s}}{1-q^{-1-s}},$$

with $N(g) = Card\left(\left\{(u,v)\in\left(\mathbb{F}_q^{\times}\right)^2 \mid g(u,v)=0\right\}\right)$.

3.7. **Computation of $Z(s,g,v,\Delta_i)$, $i=1,2,3,4,6,7,8,9$.** These integrals can be computed as the corresponding integrals in the previous example. The following table summarizes these computations.

(3.18)

Cone	$Z(s,g,v,\Delta)$
Δ_1	$q^{-1}(1-q^{-1})$
Δ_2	$(1-q^{-1})\frac{q^{-3-6s}}{(1-q^{-2-6s})}$
Δ_3	$(1-q^{-1})^2\frac{q^{-2-6s}}{(1-q^{-2-6s})}$
Δ_4	$(1-q^{-1})^2\frac{q^{-7-24s}}{(1-q^{-2-6s})(1-q^{-5-18s})}$
Δ_6	$\frac{(1-q^{-1})^2q^{-8-27s}}{(1-q^{-5-18s})(1-q^{-3-9s})}$
Δ_7	$\frac{(1-q^{-1})^2q^{-3-9s}}{(1-q^{-3-9s})}$
Δ_8	$\frac{(1-q^{-1})q^{-4-9s}}{(1-q^{-3-9s})}$
Δ_9	$q^{-1}(1-q^{-1})$

3.8. **Computation of $Z(s,f,v,\Delta_5)$ (an integral on a degenerate face in the sense of Kouchnirenko).**

$$\begin{aligned} Z(s,g,v,\Delta_5) &= \sum_{n=1}^{\infty} \int\limits_{\pi^{3n}O_v^{\times}\times\pi^{2n}O_v^{\times}} \mid g(x,y) \mid_v^s \mid dxdy \mid \\ (3.19) = & \sum_{n=1}^{\infty} q^{-5n-18ns} \int\limits_{O_v^{\times 2}} \mid (y^3-x^2)^2(y^3-ax^2)+\pi^{2n}x^4y^4 \mid_v^s \mid dxdy \mid . \end{aligned}$$

We set $g^{(n)}(x,y) = (y^3-x^2)^2(y^3-ax^2)+\pi^{2n}x^4y^4$, for $n \geqslant 1$. In order to compute (3.19), we first compute the integral

$$I(s,g^{(n)},v) := \int\limits_{O_v^{\times 2}} \mid (y^3-x^2)^2(y^3-ax^2)+\pi^{2n}x^4y^4 \mid_v^s \mid dxdy \mid , \; n \geqslant 1. \tag{3.20}$$

By using the change of variables $\Phi(x,y)$ defined in (3.6), we have that

$$g^{(n)}\circ\Phi(x,y) = x^{18}y^6\widetilde{g^{(n)}}(x,y), \tag{3.21}$$

with

$$\widetilde{g^{(n)}}(x,y)=(y-1)^2(y-a)+\pi^{2n}x^2y^2. \tag{3.22}$$

Since $\Phi(x,y)$ gives a bijection of $O_v^{\times 2}$ onto itself preserving the Haar measure, it follows from (3.20), and (3.21), that

$$I(s,g^{(n)},v)=\int_{O_v^{\times 2}} \mid \widetilde{g^{(n)}}(x,y)\mid_v^s \mid dxdy\mid . \tag{3.23}$$

In order to compute the integral $I(s,g^{(n)},v)$, we decompose $O_v^{\times 2}$ as follows:

$$\begin{aligned} O_v^{\times 2} &= \left(O_v^{\times}\times\{y_0+\pi O_v \mid y_0\not\equiv 1,a \bmod \pi\}\right)\bigcup\left(O_v^{\times}\times\{1+\pi O_v\}\right) \\ &\bigcup\left(O_v^{\times}\times\{a+\pi O_v\}\right). \end{aligned} \tag{3.24}$$

From the above partition follows that

$$\begin{aligned} I(s,g^{(n)},v) &= (1-q^{-1})q^{-1}(q-3) \\ &+\sum_{j=1}^{\infty} q^{-j}\int_{O_v^{\times}\times O_v^{\times}} \mid \widetilde{g^{(n)}}(x,1+\pi^j y)\mid_v^s \mid dxdy\mid \\ &+\sum_{j=1}^{\infty} q^{-j}\int_{O_v^{\times}\times O_v^{\times}} \mid \widetilde{g^{(n)}}(x,a+\pi^j y)\mid_v^s \mid dxdy\mid, \end{aligned} \tag{3.25}$$

with

$$\widetilde{g^{(n)}}(x,1+\pi^j y)=\pi^{2j}\gamma_{1,j}(x,y)y^2+\pi^{2n}\gamma_{2,j}(x,y)x^2, \tag{3.26}$$

$$\gamma_{1,j}(x,y)=1-a+\pi^j y,\quad \gamma_{2,j}(x,y)=(1+\pi^j y)^2, \tag{3.27}$$

and

$$\widetilde{g^{(n)}}(x,a+\pi^j y)=\pi^{j}\delta_{1}(x,y)y+\pi^{2n}\delta_{2}(x,y)x^2, \tag{3.28}$$

$$\delta_{1,j}(x,y)=(a-1+\pi^j y)^2,\quad \delta_{2,j}(x,y)=(a+\pi^j y)^2.$$

In the next step, we compute explicitly $\mid \widetilde{g^{(n)}}(x,1+\pi^j y)\mid_v^s$, and $\mid \widetilde{g^{(n)}}(x,a+\pi^j y)\mid_v^s$, for a fixed $n\geqq 1$. By some simple calculations, it holds that

(3.29)

$$\mid \widetilde{g^{(n)}}(x,1+\pi^j y)\mid_v^s=\begin{cases} q^{-2js} & \text{if } \ 1\leqq j\leqq n-1, \\ q^{-2ns}\mid \gamma_{1,j}(x,y)y^2+\gamma_{2,j}(x,y)x^2\mid_v^s & \text{if } \ j=n, \\ q^{-2ns} & \text{if } \ j\geqq n+1, \end{cases}$$

for any $(x,z)\in O_v^{\times 2}$, and

$$
| \widetilde{g^{(n)}}(x,a+\pi^j y) |_v^s = \begin{cases} q^{-js} & \text{if } \ 1\leqq j\leqq 2n-1, \\ q^{-2ns} \mid \delta_{1,j}(x,y)y+\delta_{2,j}(x,y)x^2 \mid_v^s & \text{if } \ j=2n, \\ q^{-2ns} & \text{if } \ j\geqq 2n+1. \end{cases} \tag{3.30}
$$

We note that the explicit computation of the absolute values of $| \widetilde{g^{(n)}}(x,1+\pi^j y) |_v$, respectively, $| \widetilde{g^{(n)}}(x,a+\pi^j y) |_v$ depends on an explicit description of the set

$$
\left\{(w,z)\in\mathbb{R}^2 \mid w\leqq \min\{2z,2n\}\right\},
$$

respectively, of the set

$$
\left\{(w,z)\in\mathbb{R}^2 \mid w\leqq \min\{z,2n\}\right\},
$$

for a fixed n.

Now, from (3.25), (3.29), (3.30) and the identity $\sum\limits_{k=A}^{B} z^k=\frac{z^A-z^{B+1}}{1-z}$, we obtain the following explicit expression for $I(s,g^{(n)},v)$:

$$
\begin{aligned}
I(s,g^{(n)},v) \ = \ & q^{-1}(1-q^{-1})(q-3)+\left(1-q^{-1}\right)^2\frac{q^{-1-2s}}{1-q^{-1-2s}} \\
& -\left(1-q^{-1}\right)^2\frac{q^{-n-2ns}}{1-q^{-1-2s}}+q^{-n-2ns}I_1(s)+q^{-1}\left(1-q^{-1}\right)q^{-n-2ns} \\
& +\left(1-q^{-1}\right)^2\frac{q^{-1-s}}{1-q^{-1-s}}-\left(1-q^{-1}\right)^2\frac{q^{-2n-2ns}}{1-q^{-1-s}}+q^{-2n-2ns}I_2(s) \\
& +q^{-1}\left(1-q^{-1}\right)q^{-2n-2ns},
\end{aligned} \tag{3.31}
$$

where

$$
\begin{aligned}
I_1(s) \ : & = \int\limits_{O_v^{\times 2}} \mid \gamma_{1,n}(x,y)y^2+\gamma_{2,n}(x,y)x^2 \mid_v^s \mid dxdz \mid, \\
I_2(s) \ : & = \int\limits_{O_v^{\times 2}} \mid \delta_{1,2n}(x,y)y+\delta_{2,2n}(x,y)x^2 \mid_v^s \mid dxdz \mid .
\end{aligned} \tag{3.32}
$$

Since the reduction modulo π of the polynomials $\gamma_{1,n}(x,y)y^2+\gamma_{2,n}(x,y)x^2$ and $\delta_{1,2n}(x,y)y+\delta_{2,2n}(x,y)x^2$ does not have singular points on $\mathbb{F}_q^{\times 2}$, because the characteristic of the residue field is different from 2, SPF implies that

$$
\begin{aligned}
I_1(s) & = \frac{U_1(q^{-s})}{1-q^{-1-s}}, \\
I_2(s) & = \frac{U_2(q^{-s})}{1-q^{-1-s}},
\end{aligned} \tag{3.33}
$$

where $U_1(q^{-s})$, and $U_1(q^{-s})$ are polynomials in q^{-s}, independent of j, and n.

Finally, from (3.19), and (3.31), we obtain the following explicit expression for $Z(s,f,v,\Delta_5)$:

$$
\begin{aligned}
Z(s,f,v,\Delta_5) \;=\; & \frac{(q-3)(1-q^{-1})q^{-6-18s}}{(1-q^{-5-18s})}+\frac{(1-q^{-1})^2q^{-6-20s}}{(1-q^{-1-2s})(1-q^{-5-18s})}\\
&-\frac{(1-q^{-1})^2q^{-6-20s}}{(1-q^{-1-2s})(1-q^{-6-20s})}+\frac{q^{-6-20s}U_1(q^{-s})}{(1-q^{-1-s})(1-q^{-6-20s})}\\
&+\frac{(1-q^{-1})q^{-7-20s}}{(1-q^{-6-20s})}+\frac{(1-q^{-1})^2q^{-6-19s}}{(1-q^{-1-s})(1-q^{-5-18s})}\\
&-\frac{(1-q^{-1})^2q^{-7-20s}}{(1-q^{-1-s})(1-q^{-7-20s})}+\frac{q^{-7-20s}U_2(q^{-s})}{(1-q^{-1-s})(1-q^{-7-20s})}\\
&+\frac{(1-q^{-1})q^{-8-20s}}{(1-q^{-7-20s})}. \qquad (3.34)
\end{aligned}
$$

From the explicit formulas for $Z(s,f,v,\Delta_i)$, $i=1,2,\cdots,8,$ and $Z(s,f,v,O_v^{\times 2})$ (see (3.18), (3.34)), it follows that the real parts of the poles of $Z(s,f,v)$ belong to the set

$$\{-1\}\bigcup\{-\frac{5}{18},-\frac{1}{3}\}\bigcup\{-\frac{1}{2},-\frac{3}{10}\}\bigcup\{-\frac{7}{20}\}. \qquad (3.35)$$

4. ARITHMETIC NEWTON POLYGONS AND ARITHMETIC NON DEGENERATE POLYNOMIALS

In this section we introduce the notions of arithmetic Newton polygon and arithmetic non degeneracy for polynomials in two variables. The set of arithmetically non degenerate polynomials contains strictly the set of non degenerate polynomials in the sense of Kouchnirenko. In addition, an explicit list of candidates for the poles of a local zeta function associated to an arithmetically non degenerate polynomial can be given in terms of the corresponding arithmetic Newton polygon (see section 6).

4.1. **Arithmetically non degenerate semi-quasihomogeneous polynomials.**

Definition 4.1. *Let K be a field, and a, b two coprime positive integers. A polynomial $g(x,y)\in K[x,y]$ is quasihomogeneous with respect to the weight (a,b), if it has the form*

$$g(x,y)=cx^uy^v\prod_{i=1}^{l}\left(y^a-\alpha_ix^b\right)^{e_i},\;\; c\in K^{\times}. \qquad (4.1)$$

From the definition follows that a quasihomogeneous polynomial $g(x,y)$ satisfies

$$g(t^ax,t^by)=t^dg(x,y),\text{ for every }\; t\in K^{\times}. \qquad (4.2)$$

The integer d is called the *weighted degree* of $g(x,y)$ with respect to (a,b). We note that our definition of quasihomogeneity coincides with the standard one (i.e. with (4.2)) after a finite extension of K.

Definition 4.2. *A polynomial $g(x,y)\in K[x,y]$ is a semi-quasihomogeneous polynomial with respect to the weight (a,b), if it has the form*

$$g(x,y)=\sum_{j=0}^{l_g} g_j(x,y), \tag{4.3}$$

where each $g_j(x,y)$ is quasihomogeneous of weighted degree d_j with respect to (a,b), and $d_0<d_1<\cdots<d_{l_g}$.

The polynomial $g_0(x,y)$ is called *the quasihomogeneous part* of $g(x,y)$. If

$$g_0(x,y)=cx^{u_0}y^{v_0}\prod_{i=1}^{l_0}\left(y^a-\alpha_{i,0}x^b\right)^{e_{i,0}}$$

has singular points on the torus $\left(K^\times\right)^2$, i.e. if $e_{i,0}>1$, for some $i=1,2,\cdots,l_0$, then $g(x,y)$ is degenerate, in the sense of Kouchnirenko, with respect to its geometric Newton polygon $\Gamma^{geom}(g)$.

We put

$$f_j(x,y):=c_jx^{u_j}y^{v_j}\prod_{i=1}^{l_j}\left(y^a-\alpha_{i,j}x^b\right)^{e_{i,j}}, \ \ c_j\in K^\times, \tag{4.4}$$

d_j for the weighted degree of $f_j(x,y)$ with respect to (a,b), i.e.

$$d_j:=ab\left(\sum_{i=1}^{l_j}e_{i,j}\right)+au_j+bv_j,$$

and

$$f(x,y)=\sum_{j=0}^{l_f} f_j(x,y), \tag{4.5}$$

with $d_0<d_1<\cdots<d_{l_f}$. From now on, we shall suppose that $f(x,y)$ is a degenerate polynomial in the sense of Kouchnirenko.

Definition 4.3. *Let $f(x,y)\in K[x,y]$ be a semi-quasihomogeneous polynomial, $\theta\in K^\times$ a fixed root of $f_0(1,\theta^a)=0$, and $e_{j,\theta}$ the multiplicity of θ as root of the polynomial function $f_j(1,y^a)$. To each $f_j(x,y)$ in (4.5), we associate an straight line of the form*

$$w_{j,\theta}(z):=(d_j-d_0)+e_{j,\theta}z, \ j=0,1,\cdots,l_f,$$

and define the arithmetic Newton polygon $\Gamma_{f,\theta}$ of $f(x,y)$ at θ as

$$\Gamma_{f,\theta}:=\{(z,w)\in\mathbb{R}_+^2 \mid w\leqq\min_{0\le j\le l_f}\{w_{j,\theta}(z)\}.$$

The arithmetic Newton polygon $\Gamma^{A}(f)$ *of* $f(x,y)$ *is defined to be the set*

$$\Gamma^{A}(f) := \{\Gamma_{f,\theta} \mid \theta \in K^{\times},\ \ f_0(1,\theta^{a}) = 0\}.$$

We note that θ is one of the $\alpha_{i,0}$. In addition, $\Gamma^{A}(f)$ is a finite collection of convex sets, and no just a convex set as is $\Gamma^{geom}(f)$.

If the polynomial $f_0(x,y)$ is interpreted as the weighted tangent cone of $f(x,y)$, then for each tangent direction of the form $(\theta : 1)$, $\theta \in K^{\times}$, there exists an arithmetic Newton polygon $\Gamma_{f,\theta}$ at θ, and all these Newton polygons constitute the arithmetic Newton polygon $\Gamma^{A}(f)$ of $f(x,y)$.

Let θ be a fixed root of $f_0(1,\theta^{a}) = 0$. A point Q of the topological boundary of $\Gamma_{f,\theta}$ is called a *vertex*, if $Q = (0,0)$ or if it is the intersection point of two different straight lines of the form $w_{j,\theta}(z)$. Let Q_k, $k = 0,1,\cdots,r$, denote the vertices of the topological boundary of $\Gamma_{f,\theta}$, with $Q_0 := (0,0)$. The boundary of $\Gamma_{f,\theta}$ is conformed by r straight segments, a half-line, and the non-negative part of the horizontal axis of the $(w,z)-$plane. We put

$$w_{k,\theta}(z) = (\mathcal{D}_k - d_0) + \mathcal{E}_k z, \quad k = 1,2,\cdots,r, \tag{4.6}$$

for the equation of the straight segment between Q_{k-1} and Q_k, and

$$w_{r+1,\theta}(z) = (\mathcal{D}_{r+1} - d_0) + \mathcal{E}_{r+1} z, \tag{4.7}$$

for the equation of the half-line starting at Q_r. With this notation it has that

$$Q_k = (\tau_k, (\mathcal{D}_k - d_0) + \mathcal{E}_k \tau_k),\ k = 1,2,\cdots,r, \tag{4.8}$$

with

$$\tau_k := \frac{(\mathcal{D}_{k+1} - \mathcal{D}_k)}{\mathcal{E}_k - \mathcal{E}_{k+1}} > 0,\ k = 1,2,\cdots,r. \tag{4.9}$$

We note that $\mathcal{D}_k = d_{j_k}$, and $\mathcal{E}_k = e_{j_k,\theta}$, for some index j_k. In particular, $\mathcal{D}_1 = d_0$, $\mathcal{E}_1 = e_{0,\theta}$, and the first equation is $w_{1,\theta}(z) = \mathcal{E}_1 z$. The slope $\mathcal{E}_{r+1}$ may be 0.

Let Q be a vertex of the boundary of $\Gamma_{f,\theta}$. *The face function* $f_Q(x,y)$ is defined to be the polynomial

$$f_Q(x,y) := \sum_{w_{j,\theta}(Q)=0} f_j(x,y), \tag{4.10}$$

where $w_{j,\theta}(z)$ is the straight line corresponding to $f_j(x,y)$.

Now we introduce a new notion of non degeneracy with respect to an arithmetic Newton polygon.

Definition 4.4. *A semi-quasihomogeneous polynomial* $f(x,y) \in K[x,y]$ *is called arithmetically non degenerate with respect to* $\Gamma_{f,\theta}$ *at* θ, *if it satisfies:*

(1) *the origin of* K^2 *is a singular point of* $f(x,y)$;

(2) *the polynomial* $f(x,y)$ *does not have singular points on* $(K^{\times})^2$;
(3) *the system of equations*

$$f_Q(x,y)=\frac{\partial f_Q}{\partial x}(x,y)=\frac{\partial f_Q}{\partial y}(x,y)=0$$

does not have solutions on $(K^{\times})^2$, *for each vertex* $Q\neq(0,0)$ *of boundary of* $\Gamma_{f,\theta}$.

A semi-quasihomogeneous polynomial $f(x,y)\in K[x,y]$ *is called arithmetically non degenerate with respect to* $\Gamma^A(f)$, *if it is arithmetically non degenerate with respect to* $\Gamma_{f,\theta}$, *for each* $\theta\in K^{\times}$ *satisfying* $f_0(1,\theta^a)=0$.

4.2. **Arithmetically non degenerate polynomials.** Let $f(x,y)\in K[x,y]$ be a non-constant polynomial, and $\Gamma^{geom}(f)$ its geometric Newton polygon. Each facet $\gamma\subset\Gamma^{geom}(f)$ has a perpendicular primitive vector $a_\gamma=(a_1(\gamma),a_2(\gamma))$ and the corresponding supporting line has an equation of the form $\langle a_\gamma,x\rangle=d(\gamma)$.

Definition 4.5. *Let* $f(x,y)\in K[x,y]$ *be a non-constant polynomial such that* $f(x,y)$ *is semi-quasihomogeneous with respect to the weight* a_γ, *in the sense of the definition 4.2, for every facet* γ *of* $\Gamma^{geom}(f)$. *Let* $\Gamma^A(f,\gamma)$ *be the arithmetic Newton polygon of* $f(x,y)$ *considered as a semi-quasihomogeneous polynomial with respect to the weight* a_γ. *The arithmetic Newton polygon* $\Gamma^A(f)$ *of* $f(x,y)$ *is defined to be the set*

$$\Gamma^A(f):=\bigcup_{\{\gamma\ a\ facet\ of\ \Gamma^{geom}(f)\}}\Gamma^A(f,\gamma).$$

We note that the construction of the arithmetic polygon of f needs that f_γ can be factored as in (4.4), for every facet γ of $\Gamma^{geom}(f)$. This condition can always be attained by passing to a finite extension of K.

Definition 4.6. *A polynomial* $f(x,y)\in K[x,y]$ *is called arithmetically non degenerate with respect to its arithmetic Newton polygon, if for every facet* γ *of* $\Gamma^{geom}(f)$, *the semi-quasihomogeneous polynomial* $f(x,y)$, *with respect to the weight* a_γ, *is arithmetically non degenerate with respect to* $\Gamma^A(f,\gamma)$.

We put $K=\mathbb{C}$, and fix a geometric Newton polygon $\Gamma^{geom}\subset\mathbb{R}^2$. Then the set of degenerate polynomials in two variables with respect Γ^{geom} contains an open set, for the Zariski topology, consisting of arithmetically non degenerate polynomials. The proof of this fact is similar to [1, page 157].

4.3. **Examples.** The following three examples illustrate the above definitions for specific polynomials, we assume that K is a field of characteristic zero.

Example 4.1. *We set* $f(x,y) = (y^3 - x^2)^2 + x^4y^4 \in K[x,y]$. *The polynomial* $f(x,y)$ *is a degenerate, in the sense of Kouchnirenko, semi-quasihomogeneous polynomial with respect to the weight* $(3,2)$. *The origin of* K^2 *is the only singular point of* $f(x,y)$. *In this case the arithmetic Newton polygon of* $f(x,y)$ *is equal to* $\Gamma_{f,1}$. *The boundary of the Newton polygon* $\Gamma_{f,1}$ *is conformed by the straight segments*

$$\begin{aligned} w_{1,1}(z) &= 2z, \quad 0 \le z \le 4, \\ w_{2,1}(z) &= 8, \quad z \geqslant 4, \end{aligned}$$

and the half-line $\left\{(z,w) \in \mathbb{R}_+^2 \mid w = 0\right\}$. *Thus* $\mathcal{D}_1 = d_0 = 12$, $\mathcal{E}_1 = 2$,$\mathcal{D}_2 = 20$, $\tau_0 = 0$, $\tau_1 = 4$. *The arithmetic Newton polygon* $\Gamma_{f,1}$ *is showed in the next figure.*



FIGURE 1. $\Gamma^A(f)$

The face functions are

$$f_{(0,0)}(x,y) = (y^3 - x^2)^2, \;\; f_{(4,8)}(x,y) = (y^3 - x^2)^2 + x^4y^4.$$

Since $f_{(4,8)}(x,y)$ *does not have singular points on* $K^{\times 2}$, $f(x,y)$ *is arithmetically non degenerate.*

Example 4.2. *We set*

$$g(x,y) = (y^3 - x^2)^5 + (y^3 - x^2)^3x^6y^3 + (y^3 - x^2)^2x^{12} + x^{24} \in K[x,y].$$

The polynomial $g(x,y)$ *is a degenerate, in the sense of Kouchnirenko, semi-quasihomogeneous polynomial with respect to the weight* $(3,2)$. *The origin of* K^2 *is the only singular point of* $g(x,y)$. *It can be decomposed as*

$$g(x,y) = g_0(x,y) + g_1(x,y) + g_2(x,y) + g_3(x,y),$$

where

$$\begin{aligned} g_0(x,y) &= (y^3 - x^2)^5, \text{ hence } d_0 = 30, \; e_{0,1} = 5 \text{ and } w_{0,1} = 5z; \\ g_1(x,y) &= (y^3 - x^2)^3x^6y^3, \text{ hence } d_1 = 42, \; e_{1,1} = 3 \text{ and } w_{1,1} = 12 + 3z; \\ g_2(x,y) &= (y^3 - x^2)^2x^{12}, \text{ hence } d_2 = 48, \; e_{2,1} = 2 \text{ and } w_{2,1} = 18 + 2z; \\ g_3(x,y) &= x^{24}, \text{ hence } d_3 = 72, \; e_{3,1} = 0 \text{ and } w_{3,1} = 42. \end{aligned}$$

Since $g_0(1,\theta^a)=0$ *has only one root,* $\theta=1$*, the arithmetic Newton polygon* $\Gamma^A(g)$ *of* $g(x,y)$ *is equal to* $\Gamma_{g,1}$*. The boundary of* $\Gamma^A(g)$ *is conformed by the straight segments*

$$\begin{aligned} w_{1,1}(z) &= 5z, \quad 0\le z\le 6, \\ w_{2,1}(z) &= 18+2z, \quad 6\le z\le 12, \\ w_{3,1}(z) &= 42, \quad z\geqslant 12, \end{aligned}$$

and the half-line $\{(z,w)\in\mathbb{R}_+^2 \mid w=0\}$*. Thus* $\mathcal{D}_1=d_0=30$*,* $\mathcal{E}_1=5$*,* $\mathcal{D}_2=48$*,* $\mathcal{E}_2=2$*,* $\mathcal{D}_3=72$*,* $\mathcal{E}_3=0$*,* $\tau_0=0$*,* $\tau_1=6$*,* $\tau_2=12$*.*

We observe that the line $w(z)=12+3z$ *meets* $\Gamma_{g,1}$ *only at* $(6,30)$*. The arithmetic Newton polygon of* g *is showed in the next figure.*



FIGURE 2. $\Gamma^A(g)$.

The face functions are

$$\begin{aligned} g_{(0,0)}(x,y) &= (y^3-x^2)^5, \\ g_{(6,30)}(x,y) &= (y^3-x^2)^5+(y^3-x^2)^3x^6y^3+(y^3-x^2)^2x^{12}, \\ g_{(12,40)}(x,y) &= (y^3-x^2)^2x^{12}+x^{24}. \end{aligned}$$

Since $g_{(6,30)}(x,y)=(y^3-x^2)^2\left((y^3-x^2)^3+(y^3-x^2)+x^{12}\right)$ *has singular points on* $(K^\times)^2$*,* $g(x,y)$ *is arithmetically degenerate.*

Example 4.3. *We set*

$$h(x,y)=(y^5-x^3)^4(y^5-ax^3)(y^5-bx^3)+x^{20}\in K[x,y],$$

with $1\neq a\neq b$*. The polynomial* $h(x,y)$ *is a degenerate, in the sense of Kouchnirenko, semi-quasihomogeneous polynomial with respect the weight* $(5,3)$*. The origin of* K^2 *is a singular point of* $h(x,y)$*, and there are no singularities on the torus* $(K^\times)^2$*. In this case the arithmetic Newton polygon of* $h(x,y)$ *is equal to* $\Gamma^A(h)=\{\Gamma_{h,1},\Gamma_{h,a},\Gamma_{h,b}\}$*. The boundary of Newton polygon* $\Gamma_{h,1}$ *is conformed by the straight segments*

$$\begin{aligned} w_{1,1}(z) &= 4z, \quad 0\le z\le \frac{5}{2}, \\ w_{2,1}(z) &= 10, \quad z\geqslant \frac{5}{2}, \end{aligned}$$

and the half-line $\left\{(z,w)\in\mathbb{R}_+^2 \mid w=0\right\}$. *The arithmetic Newton polygon* $\Gamma_{h,1}$ *is showed in the next figure.*



FIGURE 3. $\Gamma_{h,1}$.

The face functions are

$$\begin{aligned} h_{(0,0)}(x,y) &= (y^5-x^3)^4(y^5-ax^3)(y^5-bx^3) \\ h_{(\frac{5}{2},10)}(x,y) &= (y^5-x^3)^4(y^5-ax^3)(y^5-bx^3)+x^{20}. \end{aligned}$$

Since

$$h_{(\frac{5}{2},10)}(x,y) = (y^5-x^3)^4(y^5-ax^3)(y^5-bx^3)+x^{20}$$

has no singular points on $(K^\times)$, $f(x,y)$ *is arithmetically non degenerate with respect to* $\Gamma_{h,1}$.

The boundary of the Newton polygon $\Gamma_{h,a}$ *is conformed by the straight segments*

$$\begin{aligned} w_{1,a}(z) &= z, \quad 0\le z\le 10, \\ w_{2,a}(z) &= 10, \quad z\geqslant 10, \end{aligned}$$

and the half-line $\left\{(z,w)\in\mathbb{R}_+^2 \mid w=0\right\}$. *The arithmetic Newton diagram* $\Gamma_{h,a}$ *is showed in the next figure.*



FIGURE 4. $\Gamma_{h,a}$.

The face functions are

$$\begin{aligned} h_{(0,0)}(x,y) &= (y^5-x^3)^4(y^5-ax^3)(y^5-bx^3), \\ h_{(10,10)}(x,y) &= (y^5-x^3)^4(y^5-ax^3)(y^5-bx^3)+x^{20}. \end{aligned}$$

Since

$$h_{(10,10)}(x,y) = (y^5-x^3)^4(y^5-ax^3)(y^5-bx^3)+x^{20}$$

has no singular points on $(K^{\times})^2$, $f(x,y)$ *is arithmetically non degenerate with respect to* $\Gamma_{h,a}$. *The Newton polygon* $\Gamma_{h,b}$ *is equal to* $\Gamma_{h,a}$. *Thus the polygnomial* $h(x,y)$ *is arithmetically non degenerate with respect* Γ_h^A.

5. Degenerate integrals of type $Z(s,f,v,\Delta)$

In this section using the notion of arithmetic non degeneracy introduced in the above section, we shall compute explicitly degenerate integrals of type

$$Z(s,f,v,\Delta) = \int\limits_{E} \mid f(x,y) \mid_v^s \mid dxdy \mid,$$

where

$$\Delta := (a,b)\mathbb{R}_+, \text{ and } \; E := \{(x,y) \in O_v^2 \mid (v(x),v(y)) \in \Delta\},$$

and $f(x,y) \in O_v[x,y]$ is a degenerate, in the sense of Kouchnirenko, semi-quasihomogeneous polynomial with respect to the weight (a,b), but arithmetically non degenerate with respect $\Gamma^A(f) = \cup_\theta \Gamma_{f,\theta}$. From now on, we shall consider only those $\Gamma_{f,\theta}$ for which $\theta \in O_v$. The reason is that the $\Gamma_{f,\theta}$, with $\theta \in L_v \setminus O_v$, do not contribute the poles of $Z(s,f,v,\Delta)$.

Definition 5.1. *For a semi-quasihomogeneous polynomial* $f(x,y) \in L_v[x,y]$ *non degenerate with respect to* $\Gamma^A(f) = \cup_{\{\theta\ \in O_v \mid f_0(1,\theta^a)=0\}} \Gamma_{f,\theta}$, *we define*

(5.1)
$$\mathcal{P}(\Gamma_{f,\theta}) := \bigcup_{i=1}^{r_\theta} \{-\frac{1}{\mathcal{E}_i}, -\frac{(a+b)+\tau_i}{\mathcal{D}_{i+1}+\mathcal{E}_{i+1}\tau_i}, -\frac{(a+b)+\tau_i}{\mathcal{D}_i+\mathcal{E}_i\tau_i}\} \bigcup \bigcup_{\{\mathcal{E}_{r+1}\neq o\}} \{-\frac{1}{\mathcal{E}_{r+1}}\},$$

and

$$\mathcal{P}(\Gamma^A(f)) := \bigcup_{\{\theta\ \in O_v \mid f_0(1,\theta^a)=0\}} \mathcal{P}(\Gamma_{f,\theta}). \tag{5.2}$$

The data $\mathcal{D}_i$, $\mathcal{E}_i$, τ_i, for each i, are obtained from the equations of the straight segments conforming the boundary of $\Gamma_{f,\theta}$.

The main result of this section is the following.

Theorem 5.1. *Let* $f(x,y)=\sum_{j=0}^{l_f} f_j(x,y)\in O_v[x,y]$ *be a semi-quasihomogeneous polynomial, with respect to the weight* (a,b)*, with* a*,* b *coprime,* $f_j(x,y)$ *as in (4.4). If* $f(x,y)$ *is arithmetically non degenerate with respect to* $\Gamma^A(f)$*, then the real parts of the poles of* $Z(s,f,v,\Delta)$ *belong to the set*

$$\{-1\}\bigcup\{-\frac{a+b}{d_0}\}\bigcup\mathcal{P}(\Gamma^A(f)).$$

The proof will be given at the end of this section, and it will be accomplished by computing explicitly several $p-$adic integrals. Moreover the proof provides an effective method for computing these integrals.

The following two examples illustrate the theorem.

Example 5.1. *We set* $f(x,y)=(y^3-x^2)^2+x^4y^4\in L_v[x,y]$*. We suppose that the characteristic of the residue field is different from* 2*. The polynomial* $f(x,y)$ *is arithmetically non degenerate, semi-quasihomogeneous polynomial with respect to the weight* $(3,2)$ *(see example (4.1). In this case, we have the following data attached the arithmetic Newton polygon* $\Gamma^A(f)=\Gamma_{f,1}$*,* $a=3$*,* $b=2$*,*$\mathcal{D}_1=d_0=12$*,* $\tau_1=4,\mathcal{E}_1=2$*,* $\mathcal{D}_2=20$*. Thus according the above theorem, the real parts of the poles of the integral* $Z(s,f,v)$ *belong to the set*

$$\{-1\}\bigcup\{-\frac{5}{12}\}\bigcup\{-\frac{1}{2},-\frac{9}{20}\}.$$

In subsection (3.1), we computed explicitly the local zeta function $Z(s,f,v)$ (cf. (3.17)).

Example 5.2. *We set* $g(x,y)=(y^3-x^2)^2(y^3-ax^2)+x^4y^4\in L_v[x,y]$*, with* $a\not\equiv 1 \bmod \pi$*. We assume that the characteristic of the residue field of* L_v *is different from* 2*. The polynomial* $g(x,y)$ *is quasihomogeneous with respect to the weight* $(3,2)$*. The origin of* L_v^2 *is the only singular point of* $g(x,y)$*.*

The arithmetic Newton polygon $\Gamma^A(g)=\{\Gamma_{g,1},\ \Gamma_{g,a}\}$*. The boundary of the Newton polygon* $\Gamma_{g,1}$ *is conformed by*

$$\begin{aligned} w_{1,1}(z)&=2z,\quad 0\le z\le 1,\\ w_{2,1}(z)&=2,\quad z\geqslant 1,\\ w(z)&=0,\quad z\geqslant 0.\end{aligned}$$

Thus $\mathcal{D}_1=d_0=18$*,* $\mathcal{E}_1=2$*,* $\mathcal{D}_2=20$*,* $\mathcal{E}_2=0$*,* $\tau_0=0$*,* $\tau_1=1$*.The polynomial is arithmetically non degenerate with respect* $\Gamma_{g,1}$*. According to the previous theorem the contribution of the arithmetic Newton polygon* $\Gamma_{g,1}$ *to the set of real parts of* $Z(s,g,v)$ *is* $\{-\frac{1}{2},-\frac{3}{10}\}$*.*

The boundary of the Newton polygon $\Gamma_{g,a}$ *is conformed by*

$$\begin{aligned} w_{1,1}(z)&=z,\quad 0\le z\le 2,\\ w_{2,1}(z)&=2,\quad z\geqslant 2.\\ w(z)&=0,\quad z\geqslant 0.\end{aligned}$$

Thus $\mathcal{D}_1 = d_0 = 18$, $\mathcal{E}_1 = 1$, $\mathcal{D}_2 = 20$, $\mathcal{E}_2 = 0$, $\tau_0 = 0$, $\tau_1 = 2$. *The polynomial is arithmetically non degenerate with respect* $\Gamma_{g,a}$. *According to the previous theorem the contribution of the arithmetic Newton polygon* $\Gamma_{g,a}$ *to the set of real parts of* $Z(s, g, v)$ *is* $\{-1, -\frac{7}{20}\}$. *Then according to the previous theorem the real parts of the poles of* $Z(s, g, v)$ *belong to the set* $\{-1\}\bigcup\{-\frac{5}{18}\}\bigcup\{-\frac{1}{2}, -\frac{9}{20}\}\bigcup\{-\frac{7}{20}\}$. *In subsection (3.5), we computed explicitly the local zeta* $Z(s, g, v)$ *(cf. (3.35)).*

5.1. **Some $p-$adic integrals.**

Proposition 5.1. *Let* $f(x, y) \in O_v[x, y]$ *be a semi-quasihomogeneous polynomial, with respect to the weight* (a, b), *with* a, b *coprime, and*

$$f^{(m)}(x, y) := \pi^{-d_0 m} f(\pi^{am}x, \pi^{bm}y) = \sum_{j=0}^{l_f} \pi^{(d_j - d_0)m} f_j(x, y), \tag{5.3}$$

with $f_j(x, y)$ *as in (4.4), and* $m \geqq 1$. *Then there exists a measure-preserving bijection*

$$\Phi : O_v^{\times\ 2} \to O_v^{\times\ 2}$$
$$(u, w) \to (\Phi_1(x, y), \Phi_2(x, y)),$$

such that $f^{(m)} \circ \Phi(x, y) = u^{N_i} w^{M_i} \widetilde{f^{(m)}}(u, w)$, *with* $\widetilde{f^{(m)}}(u, w)$ *non vanishing identically on the hypersurface* $uw = 0$, *and*

$$\widetilde{f^{(m)}}(u, w) = \sum_{j=0}^{l_f} \pi^{(d_j - d_0)m} \widetilde{f}_j(u, w), \tag{5.4}$$

with

$$\widetilde{f}_j(u, w) = c_j u^{A_j} w^{B_j} \prod_{i=1}^{l_j} (w - \alpha_{i,j})^{e_{i,j}}, \quad or$$

$$\widetilde{f}_j(u, w) = c_j u^{L_j} w^{F_j} \prod_{i=1}^{l_j} (1 - \alpha_{i,j} u)^{e_{i,j}}. \tag{5.5}$$

Proof. We denote by Φ_1 the map

$$\Phi_1(u, w) : O_v^{\times\ 2} \to O_v^{\times\ 2}$$
$$(u, w) \to (x, y),$$

with $x = u$, $y = uw$, and by Φ_2 the map

$$\Phi_2(u, w) : O_v^{\times\ 2} \to O_v^{\times\ 2}$$
$$(u, w) \to (x, y),$$

with $x = uw$, $y = w$.

Since $\mid \det \Phi_1'(u, w) \mid_v = 1$, and$\mid \det \Phi_2'(u, w) \mid_v = 1$, the maps Φ_1, Φ_2 give a measure-preserving bijection of $O_v^{\times\ 2}$ to itself. We shall show that Φ is a finite composition of maps of the form Φ_1 or Φ_2. The proof will be accomplished by induction on $\min\{a, b\}$.

Case $\min\{a,b\}=1$

In this case, it is sufficient to take Φ as follows:

$$\Phi=\begin{cases}\underbrace{\Phi_1\circ\cdots\circ\Phi_1}_{b-\text{times}} & \text{if} \quad a=1,\\ \underbrace{\Phi_2\circ\cdots\circ\Phi_2}_{a-\text{times}} & \text{if} \quad b=1.\end{cases}$$

Induction hypothesis

Suppose by induction hypothesis that the proposition is valid for all polynomials $f^{(m)}(x,y)$ of the form (5.3) satisfying $1\leqq \min\{a,b\}\leqq k$, with $k\geqq 1$.

Case $\min\{a,b\}=k+1$, $k\geqq 2$

Let $f^{(m)}(x,y)$ be a polynomial of the form (5.3) satisfying $\min\{a,b\}=k+1$, $k\geqq 1$, and $a>b$. By applying the Euclidean algorithm to a, b, we have that

$$a=q_1b+r_1, \qquad 0\leqq r_1<b,$$

for some q_1, $r_1\in\mathbb{N}$. Because a, b are coprime, necessarily $1\leqq r_1<b$.

We set $\Psi=\underbrace{\Phi_2\circ\cdots\circ\Phi_2}_{q_1-\text{times}}$, i.e. $x=uw^{q_1}$, $y=w$, thus

$$f^{(m)}\circ\Psi(x,y)=u^{A_i}w^{B_i}f^{*(m)}(u,w)$$

with $f^{*(m)}(u,w)$ non vanishing identically on $uw=0$, and

$$f^{*(m)}(u,w)=\sum_{j=0}^{l_f}\pi^{(d_j-d_0)m}f_j^{*}(u,w), \tag{5.6}$$

with

$$f_j^{*}(u,w)=c_ju^{C_j}w^{D_j}\prod_{i=1}^{l_j}\left(w^{r_1}-\alpha_{i,j}u^{b}\right)^{e_{i,j}}.$$

Since $\min\{r_1,b\}=r_1$, and $1\leqq r_1\leqq b-1=k$, by applying the induction hypothesis applied to the polynomial $f^{*(m)}(u,v)$ in (5.6), there exists a map Θ, that is a finite composition of maps of the form Φ_1 or Φ_2, such that $\left(f^{(m)}\circ\Psi\right)\circ\Theta=f^{(m)}\circ(\Psi\circ\Theta)=u^{N_i}w^{M_i}\ \widetilde{f^{(m)}}(u,v)$, and $\widetilde{f^{(m)}}(u,v)$ has all announced properties in the proposition. Therefore, it is sufficient to take $\Phi=\Psi\circ\Theta$.

The case $b>a$ is proved in an analogous form. $\square$

Remark 5.1. *We can suppose without loss of generality that*

$$\widetilde{f^{(m)}}(u,w)=\sum_{j=0}^{l_f}\pi^{(d_j-d_0)m}\widetilde{f}_j(u,w), \tag{5.7}$$

with

$$(5.8)\qquad \widetilde{f_j}(u,w)=c_j u^{A_j} w^{B_j}\prod_{i=1}^{l_j}(w-\alpha_{i,j})^{e_{i,j}}\,,\quad \text{with } c_j,\ \alpha_{i,j}\in L_v^{\times}.$$

We define

$$I(s,f^{(m)},v,O_v^{\times 2}):=\int\limits_{O_v^{\times 2}} \mid f^{(m)}(x,y)\mid_v^s \mid dxdy \mid .$$

Proposition 5.2. *Let $f(x,y)\in O_v[x,y]$ be a semi-quasihomogeneous polynomial, with respect to the weight (a,b), with a, b coprime, and*

(5.9)

$$f^{(m)}(x,y)=\pi^{-d_0 m}f(\pi^{am}x,\pi^{bm}y)=\sum_{j=0}^{l_f}\pi^{(d_j-d_0)m}f_j(x,y),\quad m\geqq 1,$$

with $f_j(x,y)$ as in (4.4). There exists a constant M_0 such that if $m\geqslant M_0$, then

$$(5.10)\qquad \begin{aligned} I(s,f^{(m)},v,O_v^{\times 2}) &= U_0(q^{-s})\\ \sum_{\{\theta\in O_v\mid f_0(1,\theta^a)=0\}}\sum_{k=1}^{\infty}q^{-k}\int\limits_{O_v^{\times}\times O_v^{\times}} &\mid \widetilde{f^{(m)}}(x,\theta+\pi^k y)\mid_v^s\mid dxdy\mid, \end{aligned}$$

where $U_0(q^{-s})$ is a polynomial with rational coefficients. Moreover, the polynomial $U_0(q^{-s})$ and the constant M_0 can be computed effectively.

Proof. Proposition 5.1 implies that

(5.11)

$$I(s,f^{(m)},v,O_v^{\times 2})=\int\limits_{O_v^{\times 2}}\mid f^{(m)}(x,y)\mid_v^s\mid dxdy\mid=\int\limits_{O_v^{\times 2}}\mid \widetilde{f^{(m)}}(x,y)\mid_v^s\mid dxdy\mid .$$

We set

$$R(f_0):=\{\theta\in O_v\mid f_0(1,\theta^a)=0\},$$

$$l(f_0):=\max_{\substack{\theta\neq\theta'\\ \theta,\theta'\in R(f_0)}}\{v(\theta-\theta')\},$$

and

$$B(\theta)=B(l(f_0),\theta):=O_v^{\times}\times\left(\theta+\pi^{1+l(f_0)}O_v\right),\ \text{ for }\theta\in O_v,\text{with } v(\theta)\leq l(f_0).$$

With the above notation, if $\theta=\alpha_{i_0,0}$, then

$$v\left(\widetilde{f_j}\left(x,y\right)\right)=\begin{cases} v(c_j)+B_jv(\theta)+\sum\limits_{i=1}^{l_j} e_{i,j}v\left(\theta-\alpha_{i,j}\right) & \text{if } \ f_j\left(1,\theta\right)\neq 0,\\ v(c_j)+B_jv(\theta)+\sum\limits_{\substack{i=1\\ i\neq i_0}}^{l_j} e_{i,j}v\left(\theta-\alpha_{i,j}\right) & \\ +e_{i_0,j}\left(1+l\left(f_0\right)\right)+e_{i_0,j}v\left(\frac{y-\theta}{\pi^{1+l(f_0)}}\right) & \text{if } \ f_j\left(1,\theta\right)=0,\end{cases}$$

for every $(x,y)\in B\left(l\left(f_0\right),\theta\right)$. We put

$$Const\left(j,\theta\right):=\begin{cases} v(c_j)+B_jv(\theta)+\sum\limits_{i=1}^{l_j} e_{i,j}v\left(\theta-\alpha_{i,j}\right) & \text{if } \ f_j\left(1,\theta\right)\neq 0,\\ v(c_j)+B_jv(\theta)+\sum\limits_{\substack{i=1\\ i\neq i_0}}^{l_j} e_{i,j}v\left(\theta-\alpha_{i,j}\right) & \\ +e_{i_0,j}\left(1+l\left(f_0\right)\right) & \text{if } \ f_j\left(1,\theta\right)=0.\end{cases}$$

Then, if $\theta\notin R\left(f_0\right)$, and $m\geqq M_0$, with

$$M_0:=\max_{\substack{1\leq j\leq l_f\\ \theta}}\left\{Const\left(0,\theta\right)+1-Const\left(j,\theta\right)\right\},$$

it holds that

$$v\left(\widetilde{f^{(m)}}\left(x,y\right)\right)=Const\left(0,\theta\right),$$

for every $(x,y)\in B\left(l\left(f_0\right),\theta\right)$. From the above calculations follow that

$$\int\limits_{B(l(f_0),\theta)} \mid \widetilde{f^{(m)}}(x,y)\mid_v^s\mid dxdy\mid=\mid Const\left(0,\theta\right)\mid_v^s, \text{ if } \ \theta\notin R\left(f_0\right), \text{ and } m\geqq M_0. \tag{5.12}$$

By subdividing $O_v^{\times}\times O_v^{\times}$ into equivalence classes modulo $\pi^{1+l(f_0)}$, and using (5.11), and (5.12), we have that

$$I(s,f^{(m)},v,O_v^{\times 2})=U_0(q^{-s})+\sum_{\theta\in R(f_0)}\int\limits_{B(\theta)}\mid \widetilde{f^{(m)}}(x,y)\mid_v^s\mid dxdy\mid, \tag{5.13}$$

for $m\geqq M_0$, where $U_0(q^{-s})$ is a polynomial with rational coefficients. Moreover, the polynomial $U_0(q^{-s})$ and the constant M_0 can be computed effectively. $\square$

The integral $\int_{B(\theta)} \mid \widetilde{f^{(m)}}(x,y) \mid_v^s \mid dxdy \mid$, for $\theta \in R\left(f_0\right)$, admits the following expansion

$$\begin{aligned}
&\sum_{k=1}^{\infty} q^{-k} \int\limits_{O_v^\times \times O_v^\times} \mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v^s \mid dxdy \mid = \\
&\sum_{k=1}^{1+l(f_0)} q^{-k} \int\limits_{O_v^\times \times O_v^\times} \mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v^s \mid dxdy \mid \\
&+ \sum_{k=2+l(f_0)}^{\infty} q^{-k} \int\limits_{O_v^\times \times O_v^\times} \mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v^s \mid dxdy \mid = \\
&\frac{U_1(q^{-s})}{1-q^{-1-s}} + \sum_{k=2+l(f_0)}^{\infty} q^{-k} \int\limits_{O_v^\times \times O_v^\times} \mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v^s \mid dxdy \mid ,
\end{aligned} \tag{5.14}$$

(cf. lemma 2.1). We set

$$J(s,m,\theta) := \sum_{k=2+l(f_0)}^{\infty} q^{-k} \int\limits_{O_v^\times \times O_v^\times} \mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v^s \mid dxdy \mid .$$

Then from (5.13) and (5.14) follow that

$$I(s,f^{(m)},v,O_v^{\times 2}) = \frac{U(q^{-s})}{1-q^{-1-s}} + \sum_{\theta \in R(f_0)} J(s,m,\theta), \quad m \geqslant M_0, \tag{5.15}$$

where $U(q^{-s})$ is a polynomial with rational coefficients.

The polynomial $\widetilde{f^{(m)}}(x,\theta+\pi^k y)$ can expressed as

$$\widetilde{f^{(m)}}(x,\theta+\pi^k y) = \sum_{j=0}^{l_f} C(j,\theta) \pi^{(d_j-d_0)m+ke_{j,\theta}} \gamma_j(x,y) y^{e_{j,\theta}}, \tag{5.16}$$

for $k \geqq 2+l\left(f_0\right)$, with $C(j,\theta) = Const(j,\theta)c_j \in L_v^\times$, and $\left|\gamma_j(x,y)\right|_v = 1$, for every $(x,y) \in O_v^\times \times O_v^\times$.

Given a real number x, $[x]$ denotes the greatest integer less than or equal to x.

Proposition 5.3. *With the hypothesis of proposition 5.2. If $f(x,y)$ is arithmetically non degenerate with respect to $\Gamma_{f,\theta}$, then*

$$\begin{aligned} J(s,m,\theta) &= (1-q^{-1})^2\sum_{i=0}^{r-1}\frac{q^{-(1+s\mathcal{E}_{i+1})}}{1-q^{-(1+s\mathcal{E}_{i+1})}}q^{-(\mathcal{D}_{i+1}-d_0)ms-(1+s\mathcal{E}_{i+1})[m\tau_i]} \\ &\quad -(1-q^{-1})^2\sum_{i=0}^{r-1}\frac{1}{1-q^{-(1+s\mathcal{E}_{i+1})}}q^{-(\mathcal{D}_{i+1}-d_0)ms-(1+s\mathcal{E}_{i+1})[m\tau_{i+1}]} \\ &\quad +(1-q^{-1})^2\frac{q^{-(1+s\mathcal{E}_{r+1})}}{1-q^{-(1+s\mathcal{E}_{r+1})}}q^{-(\mathcal{D}_{r+1}-d_0)ms-(1+s\mathcal{E}_{r+1})[m\tau_r]} \\ &\quad +\sum_{i=1}^{r}q^{-(\mathcal{D}_i-d_0)ms-(1+s\mathcal{E}_i)[m\tau_i]}I_i(s), \end{aligned} \tag{5.17}$$

where τ_i, $i=0,1,\cdots,r$, *are the abscissas of the vertices of* $\Gamma_{f,\alpha_{j,0}}$, *and* $I_i(s)=\frac{M_i(q^{-s},m)}{1-q^{-1-s}}$, $M_i(q^{-s},m)\in\mathbb{Q}[q^{-s}]$, $i=1,2,\cdots,r$. *Moreover, there exists a constant* $c(f)$ *such that if* $m\geqq c(f)$, *then* $I_i(s)$ *does not dependent on* m.

Proof. The proof is based on an explicit computation of the $v-$adic absolute value of $\widetilde{f^{(m)}(x,\theta+\pi^k y)}$, when $x,\ y\in O_v^\times$. In order to accomplish it, we attach to $\widetilde{f^{(m)}}(x,\theta+\pi^k y)$ the convex set $\Gamma_{\widetilde{f^{(m)}}(x,\theta+\pi^k y)}$ defined as follows. We associate to each term

$$C(j,\theta)\pi^{(d_j-d_0)m+ke_{j,\theta}}\gamma_j(x,y)y^{e_{j,\theta}} \tag{5.18}$$

of $\widetilde{f^{(m)}}(x,\theta+\pi^k y)$ (see (5.16)) an straight line of the form

$$\widetilde{w}_{j,\theta}(\widetilde{z}):=(d_j-d_0)m+e_{j,\theta}\widetilde{z},\quad j=0,1,\cdots,l_f,$$

and associate to $\widetilde{f^{(m)}}(x,\theta+\pi^k y)$ the convex set

$$\Gamma_{\widetilde{f^{(m)}}(x,\theta+\pi^k y)}=\{(\widetilde{z},\widetilde{w})\in\mathbb{R}_+^2\mid\widetilde{w}\leqq\min_{0\leq j\leq l_f}\{\widetilde{w}_{j,\theta}(\widetilde{z})\}\}. \tag{5.19}$$

Now, we set

$$\begin{array}{rccc}\Omega_m: & \mathbb{R}^2 & \longrightarrow & \mathbb{R}^2 \\ & (u,w) & \longrightarrow & (\widetilde{u},\widetilde{w}),\end{array}$$

with $u=\frac{\widetilde{u}}{m}$, $z=\frac{\widetilde{u}}{m}$. Then $\Omega_m(\Gamma_{f,\theta})=\Gamma_{\widetilde{f^{(m)}}(x,\theta+\pi^k y)}$ (see definition 4.3). The homomorphism Ω_m sends the topological boundary of $\Gamma_{f,\theta}$ into the topological boundary of $\Gamma_{\widetilde{f^{(m)}}(x,\theta+\pi^j y)}$. Thus the vertices of $\Gamma_{\widetilde{f^{(m)}}(x,\theta+\pi^k y)}$ are

$$\Omega_m(Q_i):=\begin{cases}(0,0) & \text{if}\quad i=0,\\ (m\tau_i,(\mathcal{D}_i-d_0)m+m\mathcal{E}_i\tau_i) & \text{if}\ \ i=1,2,\cdots,r,\end{cases} \tag{5.20}$$

where the τ_i are the abscissas of the vertices of $\Gamma_{f^{(m)},\theta}$ (cf. (4.8), (4.9)).

If

$$m\geqq\max_{j,\theta}\{v\left(C(j,\theta)\right)\},$$

then $\mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v$ can be computed from $\Gamma_{\widetilde{f^{(m)}}(x,\theta+\pi^k y)}$, as follows (we use the notation introduced at (4.6), (4.8), and (4.9)):

If $m\tau_i < k < m\tau_{i+1}$, $i = 0, 1, \cdots, r-1$, then

(5.21)
$$\mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v = q^{-(\mathcal{D}_{i+1}-d_0)m-\mathcal{E}_{i+1}k} \ \left|C(i+1,\theta)\right|_v, \ (x,y) \in O_v^\times \times O_v^\times .$$

If $k > m\tau_r$, then

(5.22)
$$\mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v = q^{-(\mathcal{D}_{r+1}-d_0)m-\mathcal{E}_{r+1}k} \ \left|C(r+1,\theta)\right|_v, \ (x,y) \in O_v^\times \times O_v^\times .$$

If $k = m\tau_i$, $i = 1, 2, \cdots, r$, then

$$(5.23) \qquad \mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v =$$
$$q^{-(\mathcal{D}_i-d_0)m-\mathcal{E}_i k} \quad \mid \widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y) + \pi^{m(\mathcal{D}_{i+1}-\mathcal{D}_i)}\left(\cdots\right) \mid_v, \ (x,y) \in O_v^\times \times O_v^\times ,$$

with

$$\widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y) + \pi^{m(\mathcal{D}_{i+1}-\mathcal{D}_i)}\left(\cdots\right) = \widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y) +$$

$$\pi^{m(\mathcal{D}_{i+1}-\mathcal{D}_i)}\left(\text{terms with weighted degree } \geqq \mathcal{D}_{i+1}\right),$$

and

$$\widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y) = \sum_{\widetilde{w}_{i,\theta}(\Omega_m(Q_i))=0} \gamma_i(x,y) y^{e_{i,\theta}},$$

here $\widetilde{w}_{i,\theta}(\widetilde{z})$ is the straight line corresponding to the term

$$\pi^{(d_i-d_0)m+ke_{i,\theta}}\gamma_i(x,y)y^{e_{i,\theta}}.$$

In addition, we note any $k \in N$, $k \geqq 1$ satisfies only one of the following conditions:

$$[m\tau_i]+1 \le k \le [m\tau_{i+1}]-1, \quad i = 0, 1, \cdots, r-1;$$

$$k = [m\tau_i], \quad i = 0, 1, \cdots, r;$$

or

$$k \geqq [m\tau_r]+1.$$

Then by using (5.21), (5.22), (5.23), and the previous observation, it follows that

$$\begin{aligned} J(s,m,\theta) &= \sum_{k=2+l(f_0)}^{\infty} q^{-k} \int\limits_{O_v^{\times}\times O_v^{\times}} \mid \widetilde{f^{(m)}}(x,\theta+\pi^k y) \mid_v^s \mid dxdy \mid = \\ & (1-q^{-1})^2 \sum_{i=0}^{r-1} \left(q^{-(\mathcal{D}_{i+1}-d_0)ms} \sum_{j=[m\tau_i]+1}^{[m\tau_{i+1}]-1} q^{-(1+s\mathcal{E}_{i+1})j} \right) \\ & +(1-q^{-1})^2 \sum_{j=[m\tau_r]+1}^{\infty} q^{-(\mathcal{D}_{r+1}-d_0)ms-(1+s\mathcal{E}_{r+1})j} \\ & +\sum_{i=1}^{r} q^{-(\mathcal{D}_i-d_0)ms-(1+s\mathcal{E}_i)[m\tau_i]} I_i(s), \end{aligned} \tag{5.24}$$

with

$$I_i(s) := \int\limits_{O_v^{\times 2}} \mid \widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y) + \pi^{m(\mathcal{D}_{i+1}-\mathcal{D}_i)} \text{ (higher order terms) } \mid_v \mid dxdy \mid .$$

Since $\widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y)+\pi^{(\mathcal{D}_{i+1}-\mathcal{D}_i)}$ (higher order terms), and $\widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y)$ do not have singular points on $(L_v^{\times})^2$(cf. proposition 5.1), there exists a constant $c_0(f)$ such that

$$I_i(s) = \int\limits_{O_v^{\times 2}} \mid \widetilde{f^{(m)}_{\Omega_m(Q_i)}}(x,y) \mid_v \mid dxdy \mid = \frac{U_i(q^{-s})}{1-q^{-1-s}}, \quad U_i(q^{-s}) \in \mathbb{Q}[q^{-s}],\ i=1,\ldots,r, \tag{5.25}$$

for $m \geqq c_0(f)$ (cf. lemma 2.1). Finally, the result follows from (5.24), by using the algebraic identity

$$\sum_{k=A}^{B} z^k = \frac{z^A - z^{B+1}}{1-z}. \tag{5.26}$$

The constant $c(f) = \max\left\{c_0(f), \max_{j,\theta}\left\{v\left(C(j,\theta)\right)\right\}\right\}.$ □

5.2. **Some algebraic identities.** The following algebraic identities follows easily from (5.26). These identities will be used later on for the explicit computation of certain $p-$adic integrals. Let s be a complex number, and i, c_0 non-negative integers , with $c_0 \geqq 1$.

$$\begin{aligned} S_1(s,i,c_0) \quad : &= \sum_{m=c_0}^{\infty} q^{-(a+b)m-d_0ms-(\mathcal{D}_{i+1}-d_0)ms-(1+s\mathcal{E}_{i+1})[m\tau_i]} \\ &= \begin{cases} \frac{q^{-c_0(a+b)-c_0d_0s}}{1-q^{-(a+b)-d_0s}} & \text{if} \quad i=0, \\ \sum_{l=0}^{\mathcal{E}_i-\mathcal{E}_{i+1}-1} \left(\frac{q^{-B_{i,l}-A_l(\gamma_i+\delta_i s)}}{1-q^{-\gamma_i-\delta_i s}} \right) & \text{if} \quad i=1,\ldots,r, \end{cases} \end{aligned}$$

with

$\tau_i = \frac{\mathcal{D}_{i+1}-\mathcal{D}_i}{\mathcal{E}_i-\mathcal{E}_{i+1}},$
$A_l := \begin{cases} \left[\frac{c_0}{\mathcal{E}_i-\mathcal{E}_{i+1}}\right] & \text{if} \quad l \neq 0, \\ 1 & \text{if} \quad l = 0, \text{ and } \left[\frac{c_0}{\mathcal{E}_i-\mathcal{E}_{i+1}}\right] = 0, \\ \left[\frac{c_0}{\mathcal{E}_i-\mathcal{E}_{i+1}}\right] & \text{if} \quad l = 0, \text{ and } \left[\frac{c_0}{\mathcal{E}_i-\mathcal{E}_{i+1}}\right] \geqq 1, \end{cases}$
$B_{i,l} := (a+b)l + [l\tau_i] + s([l\tau_i]\mathcal{E}_{i+1} + l\mathcal{D}_{i+1}),$
$\gamma_i := (a+b)(\mathcal{E}_i - \mathcal{E}_{i+1}) + (\mathcal{D}_{i+1} - \mathcal{D}_i),$
$\delta_i := \mathcal{D}_{i+1}(\mathcal{E}_i - \mathcal{E}_{i+1}) + (\mathcal{D}_{i+1} - \mathcal{D}_i)\mathcal{E}_{i+1}.$

$$\begin{aligned} S_2(s,i,c_0) \quad : &= \sum_{m=1}^{\infty} q^{-(a+b)m-d_0ms-(\mathcal{D}_{i+1}-d_0)ms-(1+s\mathcal{E}_{i+1})[m\tau_{i+1}]} \\ &= \sum_{l=0}^{\mathcal{E}_{i+1}-\mathcal{E}_{i+2}-1} \left(\frac{q^{-G_{i,l}-H_l(\rho_i+\sigma_i s)}}{1-q^{-\rho_i-\sigma_i s}} \right), \quad i = 0, \cdots, r-1, \end{aligned}$$

with

$\tau_{i+1} = \frac{\mathcal{D}_{i+2}-\mathcal{D}_{i+1}}{\mathcal{E}_{i+1}-\mathcal{E}_{i+2}},$
$H_l := \begin{cases} \left[\frac{c_0}{\mathcal{E}_{i+1}-\mathcal{E}_{i+2}}\right] & \text{if} \quad l \neq 0, \\ 1 & \text{if} \quad l = 0, \text{ and } \left[\frac{c_0}{\mathcal{E}_{i+1}-\mathcal{E}_{i+2}}\right] = 0, \\ \left[\frac{c_0}{\mathcal{E}_{i+1}-\mathcal{E}_{i+2}}\right] & \text{if} \quad l = 0, \text{ and } \left[\frac{c_0}{\mathcal{E}_{i+1}-\mathcal{E}_{i+2}}\right] \geqq 1, \end{cases}$
$G_{i,l} := (a+b)l + [l\tau_{i+1}] + s([l\tau_{i+1}]\mathcal{E}_{i+1} + l\mathcal{D}_{i+1}),$
$\rho_i := (a+b)(\mathcal{E}_{i+1} - \mathcal{E}_{i+2}) + (\mathcal{D}_{i+2} - \mathcal{D}_{i+1}),$
$\delta_i := \mathcal{D}_{i+1}(\mathcal{E}_{i+1} - \mathcal{E}_{i+2}) + (\mathcal{D}_{i+2} - \mathcal{D}_{i+1})\mathcal{E}_{i+1}.$

Remark 5.2. *With the above notation, the real parts of the poles of* $S_1(s,i,c_0)$, $S_2(s,i,c_0)$, *belong to the set* $P(\Gamma_{f,\theta}) \cup \{-\frac{a+b}{d_0}\}$.

5.3. **Proof of Theorem 5.1.** The integral $Z(s,f,v,\Delta)$ admits the following expansion:

$$\begin{aligned} Z(s,f,v,\Delta) &= \sum_{m=1}^{\infty} \int\limits_{\pi^{am}O_v^{\times} \times \pi^{bm}O_v^{\times}} \mid f(x,y) \mid_v^s \mid dxdy \mid \\ &= \sum_{m=1}^{\infty} q^{-(a+b)m \; -d_0 \; ms} \int\limits_{O_v^{\times 2}} \mid f^{(m)}(x,y) \mid_v^s \mid dxdy \mid, \end{aligned} \tag{5.27}$$

with

$$f^{(m)}(x,y) = \pi^{-md_0} f(\pi^{am}x, \pi^{bm}y) = \sum_{j=0}^{l_f} \pi^{(d_j-d_0)m} f_j(x,y).$$

The integral

$$I(f^{(m)},s,v)=\int\limits_{O_v^{\times 2}} \mid f^{(m)}(x,y)\mid_v^s \mid dxdy\mid$$

is equal to

$$I(s,f^{(m)},v,O_v^{\times 2})=\frac{U_1(q^{-s})}{1-q^{-1-s}}+\sum_{\{\theta\in O_v^{\times}\mid f_0(1,\theta^a)=0\}} J(s,m,\theta), \tag{5.28}$$

for m big enough (cf. 5.15). Thus from (5.27), and (5.28), it follows that

$$\begin{aligned} Z(s,f,v,\Delta) &= \frac{U(q^{-s})}{1-q^{-1-s}}+ \\ & \sum_{\{\theta\in O_v^{\times}\mid f_0(1,\theta^a)=0\}}\left(\sum_{m=c(f)}^{\infty} q^{-(a+b)m\ -d_0\ ms}J(s,m,\theta)\right), \end{aligned} \tag{5.29}$$

where the constant $c(f)$ is defined in proposition 5.3. By using the explicit formula for $J(s,m,\theta)$ given in proposition 5.3 for $m\geqslant c(f)$, and the algebraic identities given in subsection 5.2, we have that

$$\begin{aligned} \sum_{m=c(f)}^{\infty} q^{-(a+b)m\ -d_0\ ms}J(s,m,\theta) &= \frac{U_\theta(q^{-s})}{(1-q^{-1-s})} \\ &+(1-q^{-1})^2\sum_{i=0}^{r-1}\frac{q^{-(s\mathcal{E}_{i+1}+1)}}{1-q^{-(s\mathcal{E}_{i+1}+1)}}S_1(s,i,c_0) \\ &-(1-q^{-1})^2\sum_{i=0}^{r-1}\frac{1}{1-q^{-(1+s\mathcal{E}_{i+1})}}S_2(s,i,c_0) \\ &+(1-q^{-1})^2\frac{q^{-(1+s\mathcal{E}_{r+1})}}{1-q^{-(1+s\mathcal{E}_{r+1})}}S_1(s,r,c_0) \\ &+\sum_{i=1}^{r} I_i(s)S_2(s,i-1,c_0), \end{aligned} \tag{5.30}$$

where $U_\theta(q^{-s})\in\mathbb{Q}\left[q^{-s}\right]$, and the data $\mathcal{E}_i$, $\mathcal{D}_i$, $i=1,2,\cdots,r$, depend on the arithmetic Newton polygon $\Gamma_{f,\theta.}$ Now by remark 5.2, the real parts of the poles of $S_1(s,i,c_0)$, $S_2(s,i,c_0)$, belong to the set $\mathcal{P}(\Gamma_{f,\theta})$. Then the real parts of the poles of

$$\sum_{\{\theta\in O_v^{\times}\mid f_0(1,\theta^a)=0\}}\left(\sum_{m=c(f)}^{\infty} q^{-(a+b)m\ -d_0\ ms}J(s,m,\theta)\right)$$

belong to the set $\left\{-\frac{a+b}{d_0}\right\}\bigcup \bigcup_{\{\theta\in O_v \mid f_0(1,\theta^a)=0\}} \mathcal{P}(\Gamma_{f,\theta})$, and from (5.29) follows that the real parts of the poles of $Z(s,f,v,\Delta)$ belong to the set

$$\{-1\}\bigcup\{-\frac{a+b}{d_0}\}\bigcup \bigcup_{\{\theta\in O_v \mid f_0(1,\theta^a)=0\}} \mathcal{P}(\Gamma_{f,\theta}).$$

6. Main result

In this section we prove the main result of this paper that gives an explicit list for the possible poles of a local zeta function attached to an arithmetically non degenerate polynomial in terms of the corresponding arithmetic Newton polygon.

6.1. **Local zeta functions for arithmetically non degenerate curves.** Let $f(x,y)\in L_v[x,y]$ be a non-constant polynomial satisfying $f(0,0)=0$, and

$$\mathbb{R}_+^2 = \{(0,0)\}\bigcup \bigcup_{\gamma\subset\Gamma^{geom}(f)} \Delta_\gamma, \tag{6.1}$$

a simplicial conical subdivision subordinated to $\Gamma^{geom}(f)$. We denote by $a_\gamma=(a_1(\gamma),a_2(\gamma))$ a perpendicular and primitive vector to facet γ of $\Gamma^{geom}(f)$, and by $< a, x >= d_a(\gamma)$ the equation of the corresponding supporting line. We also define

$$\mathcal{P}(\Gamma^{geom}(f)) := \{-\frac{a_1(\gamma)+a_2(\gamma)}{d_a(\gamma)} \mid \gamma \text{ a facet of } \Gamma^{geom}(f), \text{ with } d_a(\gamma)\neq 0\}.$$

The following is the main result of the present paper.

Theorem 6.1 (Main Theorem). *Let $f(x,y)\in L_v[x,y]$ be a non-constant polynomial. If $f(x,y)$ is arithmetically non degenerate with respect to its arithmetic Newton polygon $\Gamma^A(f)$, then the real parts of the poles of $Z(s,f,v)$ belong to the set*

$$\{-1\}\bigcup\mathcal{P}(\Gamma^{geom}(f))\bigcup\mathcal{P}(\Gamma^A(f)). \tag{6.2}$$

Proof. By taking a simplicial conical subdivision, the computation of $Z(s,f,v)$ is reduced to the computation of integrals of type $Z(s,f,v,O_v^{\times 2})$, and $Z(s,f,v,\Delta_\gamma)$, with γ a proper face of $\Gamma^{geom}(f)$ (see subsection 2.2). By lemma 2.1 the real parts of the poles of $Z(s,f,v,O_v^{\times 2})$ belong to the set $\{-1\}$. The computation of the integrals $Z(s,f,v,\Delta_\gamma)$, for γ a proper face of $\Gamma^{geom}(f)$, involves two cases, according if the semi-quasihomogeneous polynomial $f(x,y)$, with respect to $a_\gamma = (a_1(\gamma),a_2(\gamma))$, is geometrically non degenerate or not. If Δ_γ is a one-dimensional cone, and $f_\gamma(x,y)$ does not have singularities on $(L_v^\times)^2$, then the real parts of the poles of $Z(s,f,v,\Delta_\gamma)$ belong to the set

$$\{-1\}\bigcup\{-\frac{a+b}{d_\gamma}\}\subseteq\{-1\}\bigcup\mathcal{P}(\Gamma^{geom}(f)), \tag{6.3}$$

where $ax + by = d_\gamma$, $d_\gamma \neq 0$, is the equation of the supporting line of the facet γ (cf. lemma 2.2). If Δ_γ is a two-dimensional cone, $f_\gamma(x,y)$ is a monomial and then it does not have singularities on the torus $(L_v^\times)^2$, by lemma 2.2, $Z(s,f,v,\Delta_\gamma)$ is a entire function.

If Δ_γ is a one-dimensional cone, and $f_\gamma(x,y)$ has singularities on $(L_v^\times)^2$, then $f(x,y)$ is a semi-quasihomogeneous arithmetically non degenerate polynomial, and thus the real parts of the poles of $Z(s,f,v,\Delta_\gamma)$ belong to the set
(6.4)

$$\{-1\} \bigcup \{-\frac{a+b}{d_\gamma}\} \bigcup \mathcal{P}(\Gamma^A(f)) \subseteq \{-1\} \bigcup \mathcal{P}(\Gamma^{geom}(f)) \bigcup \mathcal{P}(\Gamma^A(f)),$$

(cf. theorem 5.1).Therefore the real part of the poles of $Z(s,f,v)$ belong to the set $\{-1\} \bigcup \mathcal{P}(\Gamma^{geom}(f)) \bigcup \mathcal{P}(\Gamma^A(f))$. □

REFERENCES

[1] Arnold V., Varchenko A. and Gussein-Zade S., Singularites des applications differentiables, vol. 2, Editions, Mir, 1986.

[2] Denef J., Report on Igusa's local zeta function, Seminaire Bourbaki 1990/1991 (730-744) in Asterisque 201-203 (1991), 359-386.

[3] Denef J., Poles of $p-$adic complex powers and Newton polyhedra, Nieuw archief voor wiskunde, 13 (1995), 289-295.

[4] Denef J., Hoornaert Kathleen, Newton polyhedra and Igusa local zeta function, To appear in Journal of Number Theory.

[5] Igusa Jun-Ichi, An introduction to the theory of local zeta functions, AMS/IP studies in advanced mathematics, v. 14, 2000.

[6] Igusa Jung-Ichi, A stationary phase formula for $p-$adic integrals and its applications, in Algebraic Geometry and its Applications, (1994), 175-1994, Springer-Verlag, New York.

[7] Igusa Jung-Ichi, Complex powers of irreducible algebroid curves, Geometry Today, Birkkhauser (1985), 207-230.

[8] Kempf G., Knudsen F., Mumford D., and Saint-Donat B., Toroidal Embeddings, Lecture Notes in Mathematics 339, Springer-Verlag, Berlin, 1973.

[9] Kouchnirenko A. G., Polyedres de Newton et nombres de Milnor, Invent. Math. 32 (1976), 1-31.

[10] Lichtin B., Meuser D., Poles of a local zeta function and Newton polygons, Compos. Math. 55 (1985), 313-332.

[11] Meuser D., On the poles of a local zeta function for curves, Invent. Math. 73 (1983), 445-465.

[12] Strauss L., Poles of two-variable $p-$adic complex power, Trans. Amer. Math. Soc., 27 (1983), 481-493.

[13] Varchenko A., Newton polyhedra and estimation of oscillanting integrals, Funct. Anal. Appl. 10 (1976), 175-196.

[14] Veys W., On the poles of Igusa's local zeta functions for curves, J. Lond. Math. Soc., 41 (1990), 27-32.

[15] Veys W., Poles of Igusa's local zeta function and monodromy, Bull. Soc. Math. Fr., 121 (1993), 545-598.

[16] Zuniga-Galindo W. A., Igusa's local zeta functions of semiquasihomogeneous polynomials, Trans. Amer. Math. Soc. 353, (2001), 3193-3207.

[17] Zuniga-Galindo W. A., local zeta functions and Newton polyhedra, to appear in Nagoya Math. J.
[18] Zuniga-Galindo W.A., Local zeta function for polynomial non-degenerate homogeneous mappings, preprint 2003.

Instituto de Matemática E Computaçao, Universidade de São Paulo at São Carlos, Av. do Trabalhador São -Carlense 400, CEP 13560-970, São Carlos - SP, Brasil
E-mail address: `mjsaia@icmc.sc.usp.br`

Department of Mathematics and Computer Science, Barry University, 11300 N.E. Second Avenue, Miami Shores, Florida 33161, USA
E-mail address: `wzuniga@mail.barry.edu`